

Inverse spectral problems for Sturm–Liouville operators with matrix-valued potentials

Ya V Mykytyuk and N S Trush

Lviv National University, 1 Universytets'ka st., 79602 Lviv, Ukraine

E-mail: yamykytyuk@yahoo.com, n.s.trush@gmail.com

Abstract. We give a complete description of the set of spectral data (eigenvalues and specially introduced norming constants) for Sturm–Liouville operators on the interval $[0, 1]$ with matrix-valued potentials in the Sobolev space W_2^{-1} and suggest an algorithm reconstructing the potential from the spectral data that is based on Krein's accelerant method.

AMS classification scheme numbers: Primary 34L20, Secondary 34B24, 34A55

Keywords: Sturm–Liouville operator, matrix-valued potential

1. Introduction

The main aim of this paper is to solve the inverse spectral problem for Sturm–Liouville operators on the unit interval $[0, 1]$ with matrix-valued distributional potentials in the Sobolev space W_2^{-1} . Before explaining the setting, we shall give a short overview of the known results; the reader is referred to the books [1–3] and the review paper [4] for further details.

1.1. Known results

The study of the inverse spectral problems for Sturm–Liouville and Dirac operators has a rather long history. It was initiated by the celebrated paper by Borg [5] of 1946, who proved that two spectra of a Sturm–Liouville equation determine uniquely the potential. Later Marchenko [6] showed that the potential is uniquely determined by the spectral function. In 1951, there appeared the paper by Gelfand and Levitan [7] giving an algorithm reconstructing the potential from the spectral measure, and the paper of Marchenko [8] with a detailed account of his method. Meanwhile Krein developed in a series of papers [9–12] an alternative approach to inverse problems. As is well explained by Ramm [13], each of the three methods—that of Gelfand and Levitan, Marchenko, and Krein—enlightens a different facet of the problem and is equally important to the theory.

Another important tool for the study of the inverse spectral problems was developed by Trubowitz and his coauthors Isaacson [14], Isaacson and McKean [15], Dalhberg [16],

and Pöschel [3]. It is based on analytic dependence of an individual eigenvalue and norming constant on the potential. In some cases (e.g. in the inverse problem for the perturbed harmonic oscillator) the Trubowitz method is probably the only one yielding a satisfactory result [17, 18].

The study of the matrix-valued Sturm–Liouville and Dirac operators has mainly followed the path of generalizing the methods developed in the scalar case. A solution to the inverse scattering problem for matrix Sturm–Liouville operators on the half-line was given by Newton and Jost [19], Krein [12, 20], Agranovich and Marchenko [21]. Analogous results for Dirac operators on the half-line were obtained in the papers of Levitan and Gasymov [22] and Gasymov [23], and for more general first order systems by Lesch and Malamud [24].

Inverse spectral problems for matrix-valued Sturm–Liouville and Dirac operators on a finite interval have not been discussed so extensively. Uniqueness of reconstruction was studied in the papers by Carlson [25] and Yurko [26], and Malamud [27] investigated uniqueness for various settings of inverse problems for systems of differential equations of the first order. Yurko in [26] and [28] suggested an algorithm reconstructing the potential from the spectral data, but did not give a complete description of the set of spectral data. Probably the closest to our work are the papers by Chelkak and Korotyaev [29, 30], which we next discuss in more detail.

In [29], the authors investigated uniqueness questions for operators

$$T_q = -\frac{d^2}{dx^2} + q \quad (1.1)$$

on $(0, 1)$ subject to the Dirichlet boundary conditions and with Hermitian matrix-valued potentials q in $L_1(0, 1)$ entrywise and studied isospectral transformations as well as some subtle properties of the local structure of the set of isospectral potentials. An important phenomenon was discovered there, namely, the one of the so called “forbidden” subspaces. In [30], the inverse spectral problem for operators T_q with potentials q belonging to $L_2(0, 1)$ entrywise was treated. The main theorem there gives a complete description of the set of spectral data. That result of [30] and Theorem 1.2 of the present paper are similar but they treat different classes of potentials. Also the approaches are different: while the proofs in [30] are based on the Trubowitz method and some special isospectral transformations, we use the Krein accelerant method.

1.2. Setting of the problem

Let M_r be the Banach algebra of the square $r \times r$ matrices with complex entries, which we identify with the Banach algebra of linear operators $\mathbb{C}^r \rightarrow \mathbb{C}^r$ endowed with the standard norm (see Appendix A). We shall write I for the unit element of M_r and M_r^+ for the set of all $A \in M_r$ such that $A = A^* \geq 0$. We shall use the abbreviations

$$W_p^s := W_p^s((0, 1), \mathbb{C}^r), \quad \mathbf{W}_p^s := W_p^s((0, 1), M_r) \quad (s \in \mathbb{Z}, p \geq 1)$$

for the corresponding Sobolev spaces, as well as the notations

$$L_p := L_p((0, 1), \mathbb{C}^r), \quad \mathbf{L}_p := L_p((0, 1), M_r), \quad p \geq 1,$$

$$Re \mathbf{L}_p := \{u \in \mathbf{L}_p \mid u = u^*\}, \quad Re \mathbf{W}_2^{-1} := \{u \in \mathbf{W}_2^{-1} \mid u = u^*\}.$$

Some further information on these spaces can be found in Appendix A.

For an arbitrary $\tau \in \mathbf{L}_2$, we consider the differential expression

$$\mathbf{t}_\tau(f) := - \left(\frac{d}{dx} + \tau \right) \left(\frac{d}{dx} - \tau \right) f$$

on the domain

$$D(\mathbf{t}_\tau) := \{f \in W_2^1 \mid f_\tau^{[1]} := (f' - \tau f) \in W_2^1\}.$$

The function $f_\tau^{[1]}$ is usually called the *quasi-derivative* of f .

Denote by $\mathbf{t}_{\tau,D}$ and $\mathbf{t}_{\tau,N}$ the restrictions of $\mathbf{t}_\tau$ onto the domains

$$D(\mathbf{t}_{\tau,D}) := \{f \in D(\mathbf{t}_\tau) \mid f \in W_{2,0}^1\},$$

$$D(\mathbf{t}_{\tau,N}) := \{f \in D(\mathbf{t}_\tau) \mid f_\tau^{[1]} \in W_{2,0}^1\}$$

respectively. Here and hereafter, $W_{2,0}^1 := \{f \in W_2^1 \mid f(0) = f(1) = 0\}$.

The differential expression $\mathbf{t}_{\tau,D}$ (unlike $\mathbf{t}_{\tau,N}$) can be written in the usual potential form. Namely, let us define the Miura map b via (cf. [31])

$$\mathbf{L}_2 \ni u \mapsto u' + u^2 =: b(u) \in \mathbf{W}_2^{-1}$$

and the class of (Hermitian) Miura potentials

$$\mathfrak{M} := \{q = b(u) \mid u \in \mathbf{L}_2\}, \quad (Re \mathfrak{M} := \{q = b(u) \mid u \in Re \mathbf{L}_2\}).$$

It can be shown (see Lemma 2.2) that for an arbitrary $q \in \mathfrak{M}$ and $\tau \in b^{-1}(q)$ one has

$$D(\mathbf{t}_{\tau,D}) = \{f \in W_{2,0}^1 \mid (-f'' + qf) \in L_2\}, \quad \mathbf{t}_{\tau,D}(f) = -f'' + qf,$$

where the derivative f'' and the product qf should be understood in the distributional sense. In particular,

$$\mathbf{t}_{\tau_1,D} = \mathbf{t}_{\tau_2,D} \quad \text{if} \quad b(\tau_1) = b(\tau_2). \quad (1.2)$$

For $\tau \in \mathbf{L}_2$ and $q := b(\tau)$, we consider the operators S_τ and T_q acting in L_2 via

$$\begin{aligned} S_\tau f &:= \mathbf{t}_{\tau,N}(f), & D(S_\tau) &:= D(\mathbf{t}_{\tau,N}), \\ T_q f &:= -f'' + qf, & D(T_q) &:= D(\mathbf{t}_{\tau,D}). \end{aligned}$$

In this paper we shall concentrate ourselves on the study of the spectral properties of the operators S_τ and T_q for $\tau \in Re \mathbf{L}_2$ and $q = b(\tau)$. In this case the operators S_τ and T_q are self-adjoint; moreover, $S_\tau \geq 0$ and $T_q > 0$. Their spectra $\sigma(S_\tau)$ and $\sigma(T_q)$ consist of a countably many isolated eigenvalues accumulating at $+\infty$; moreover, $\sigma(S_\tau) = \sigma(T_q) \cup \{0\}$.

Let $\tau \in \mathbf{L}_2$ and $\lambda \in \mathbb{C}$. We denote by $\varphi(\cdot, \lambda, \tau)$ and $\psi(\cdot, \lambda, \tau)$ the matrix-valued solutions of the Cauchy problems

$$- \left(\frac{d}{dx} + \tau \right) \left(\frac{d}{dx} - \tau \right) \varphi = \lambda^2 \varphi, \quad \varphi(0) = 0, \quad \varphi_\tau^{[1]}(0) = \lambda I, \quad (1.3)$$

$$- \left(\frac{d}{dx} - \tau \right) \left(\frac{d}{dx} + \tau \right) \psi = \lambda^2 \psi, \quad \psi(0) = I, \quad \psi_{-\tau}^{[1]}(0) = 0. \quad (1.4)$$

They are related to each other via

$$\left(\frac{d}{dx} - \tau\right) \varphi(\cdot, \lambda, \tau) = \lambda \psi(\cdot, \lambda, \tau), \quad \left(\frac{d}{dx} + \tau\right) \psi(\cdot, \lambda, \tau) = -\lambda \varphi(\cdot, \lambda, \tau)$$

and admit representations in the form (see Theorem 2.1)

$$\begin{aligned} \varphi(x, \lambda, \tau) &= \sin \lambda x I + \int_0^x (\sin \lambda t) K_{\tau, D}(x, t) dt, \\ \psi(x, \lambda, \tau) &= \cos \lambda x I + \int_0^x (\cos \lambda t) K_{\tau, N}(x, t) dt, \end{aligned} \tag{1.5}$$

where the matrix-valued kernels $K_{\tau, D}$ and $K_{\tau, N}$ belong to the algebra G_2^+ (see Appendix A). Equalities (1.5) determine uniquely $K_{\tau, D}$ and $K_{\tau, N}$ within the class $L_2(\Omega, M_r)$, $\Omega := \{(x, t) \mid 0 \leq t \leq x \leq 1\}$.

The function $\lambda \mapsto \varphi(1, \lambda, \tau)^{-1}$, as well as the Weyl–Titchmarsh function

$$m_\tau(\lambda) := -\varphi(1, \lambda, \tau)^{-1} \psi(1, \lambda, -\tau)$$

are meromorphic in $\mathbb{C}$; notice that $m_0(\lambda) = -\cot \lambda I$.

Let now $\tau \in \operatorname{Re} \mathbf{L}_2$ and $q = b(\tau)$. We denote by $\lambda_j(\tau)$ ($j \in \mathbb{Z}_+$) the square roots of the pairwise distinct eigenvalues of the operator S_τ labelled in increasing order, i.e., $0 = \lambda_0(\tau) < \lambda_k(\tau) < \lambda_{k+1}(\tau)$, $k \in \mathbb{N}$; then

$$\sigma(S_\tau) = \{\lambda_j^2(\tau)\}_{j=0}^\infty, \quad \sigma(T_q) = \{\lambda_j^2(\tau)\}_{j=1}^\infty.$$

The function m_τ is a matrix-valued Herglotz function (i.e., $\operatorname{Im} m_\tau(\lambda) \geq 0$ for $\operatorname{Im} \lambda > 0$), and the set $\{\pm \lambda_j(\tau)\}_{j \in \mathbb{Z}_+}$ is the set of its poles. Put by definition

$$\alpha_0(\tau) := -\frac{1}{2} \operatorname{res}_{\lambda=0} m_\tau(\lambda), \quad \alpha_j(\tau) := -\operatorname{res}_{\lambda=\lambda_j} m_\tau(\lambda), \quad j \in \mathbb{N}. \tag{1.6}$$

The matrix $\alpha_j(\tau)$ for $j \in \mathbb{Z}_+$ (resp. for $j \in \mathbb{N}$) is called the *norming constant* of the operator S_τ (resp. of the operator T_q) corresponding to the eigenvalue $\lambda_j^2(\tau)$. We note that the multiplicity of the eigenvalue $\lambda_j^2(\tau)$ of S_τ or T_q equals $\operatorname{rank} \alpha_j(\tau)$.

It turns out that a sequence $((\lambda_j(\tau), \alpha_j(\tau)))_{j \in \mathbb{N}}$ depends only on the function $q = b(\tau)$. In view of this we call the collections $\mathbf{a}_\tau = ((\lambda_j(\tau), \alpha_j(\tau)))_{j \in \mathbb{Z}_+}$ and $\mathbf{b}_q = ((\lambda_j(\tau), \alpha_j(\tau)))_{j \in \mathbb{N}}$ a sequences of *spectral data*, and the matrix-valued measures

$$\nu_\tau := \sum_{j=0}^\infty \alpha_j(\tau) \delta_{\lambda_j(\tau)}, \quad \mu_q := \sum_{j=1}^\infty \alpha_j(\tau) \delta_{\lambda_j(\tau)} \tag{1.7}$$

will be termed the *spectral measures* of the operators S_τ and T_q respectively. Here δ_λ is the Dirac delta-measure centred at the point λ . In particular, if $\tau = q = 0$, then

$$\nu_0 = \frac{1}{2} I \delta_0 + \sum_{n=1}^\infty I \delta_{\pi n}, \quad \mu_0 = \sum_{n=1}^\infty I \delta_{\pi n}. \tag{1.8}$$

Clearly, the spectral measures of the operators S_τ and T_q are related by the simple formula

$$\nu_\tau - \mu_q = \alpha_0(\tau) \delta_0, \quad \tau \in \operatorname{Re} \mathbf{L}_2, \quad q = b(\tau).$$

A simple relation exists also between the function m_τ and the measure ν_τ . Namely,

$$m_\tau(\lambda) = 2\lambda \int_0^\infty \frac{d\nu_\tau(\xi)}{\xi^2 - \lambda^2}, \quad \lambda \in \mathbb{C}.$$

The main aim of the present paper is to give a complete description of the classes $\mathfrak{A} := \{\mathfrak{a}_\tau \mid \tau \in Re \mathbf{L}_2\}$ and $\mathfrak{B} := \{\mathfrak{b}_q \mid q \in Re \mathfrak{M}\}$ of spectral data and to suggest an efficient method of reconstructing the functions τ and q from the measures ν_τ and μ_q respectively. We note that the description of the classes $\mathfrak{A}$ and $\mathfrak{B}$ is equivalent to the description of the families of measures $\mathcal{V} := \{\nu_\tau \mid \tau \in Re \mathbf{L}_2\}$ and $\mathcal{M} := \{\mu_q \mid q \in Re \mathfrak{M}\}$.

1.3. Main results

We start with characterization of the spectral data for Sturm–Liouville operators under consideration. In what follows $\mathfrak{a}$ (resp. $\mathfrak{b}$) will stand for an arbitrary sequence $((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ (resp. $((\lambda_j, \alpha_j))_{j \in \mathbb{N}}$), in which $(\lambda_j)_{j \in \mathbb{Z}_+}$ is a strictly increasing sequence of non-negative numbers and α_j are nonzero matrices in M_r^+ , and $\nu^{\mathfrak{a}}$ and $\mu^{\mathfrak{b}}$ will denote the measures given by

$$\nu^{\mathfrak{a}} := \sum_{j=0}^{\infty} \alpha_j \delta_{\lambda_j}, \quad \mu^{\mathfrak{b}} := \sum_{j=1}^{\infty} \alpha_j \delta_{\lambda_j}. \quad (1.9)$$

Next, we partition the semi-axis $[0, \infty)$ into pairwise disjoint intervals Δ_n ($n \in \mathbb{Z}_+$), viz.

$$\Delta_0 = \{0\}, \quad \Delta_1 = \left(0, \frac{3\pi}{2}\right], \quad \Delta_n = \left(\pi n - \frac{\pi}{2}, \pi n + \frac{\pi}{2}\right], \quad n > 1.$$

A complete description of the classes $\mathfrak{A}$ and $\mathfrak{B}$ is given by the following two theorems.

Theorem 1.1 *In order that a sequence $\mathfrak{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ should belong to $\mathfrak{A}$ it is necessary and sufficient that the following conditions be satisfied:*

$$(A_1) \quad \sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} |\lambda_j - \pi n|^2 < \infty, \quad \sup_{n \in \mathbb{N}} \sum_{\lambda_j \in \Delta_n} 1 < \infty, \quad \sum_{n=1}^{\infty} \|I - \sum_{\lambda_j \in \Delta_n} \alpha_j\|^2 < \infty;$$

$$(A_2) \quad \exists N_0 \in \mathbb{N} \quad \forall N \in \mathbb{N} \quad (N \geq N_0) \implies \sum_{n=1}^N \sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = Nr;$$

$$(A_3) \quad \text{the system of functions } \{d \cos \lambda_j x \mid j \in \mathbb{Z}_+, d \in \text{Ran } \alpha_j\} \text{ is complete in the space } L_2.$$

Theorem 1.2 *In order that a sequence $\mathfrak{b} = ((\lambda_j, \alpha_j))_{j \in \mathbb{N}}$ should belong to the class $\mathfrak{B}$ it is necessary and sufficient that the conditions (A_1) , (A_2) , and the following condition (A_4) be satisfied:*

$$(A_4) \quad \text{the system of functions } \{d \sin \lambda_j x \mid j \in \mathbb{N}, d \in \text{Ran } \alpha_j\} \text{ is complete in the space } L_2.$$

There exists a simple relation between the classes $\mathfrak{A}$ and $\mathfrak{B}$.

Proposition 1.3 *Let $\mathfrak{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ and $\mathfrak{b} = ((\lambda_j, \alpha_j))_{j \in \mathbb{N}}$. Then*

$$(\mathfrak{a} \in \mathfrak{A}) \iff (\mathfrak{b} \in \mathfrak{B}) \wedge (\lambda_0 = 0, \alpha_0 > 0).$$

By definition, every $\mathbf{a} \in \mathfrak{A}$ (resp., every $\mathbf{b} \in \mathfrak{B}$) forms spectral data for an operator S_τ with some $\tau \in \text{Re } \mathbf{L}_2$ (resp., for an operator T_q with some $q \in \text{Re } \mathfrak{M}$). It turns out that these spectral data determine the matrix-valued functions τ and q uniquely, i.e., the following holds true.

Theorem 1.4 *The mappings $\text{Re } \mathbf{L}_2 \ni \tau \mapsto \mathbf{a}_\tau \in \mathfrak{A}$ and $\text{Re } \mathfrak{M} \ni q \mapsto \mathbf{b}_q \in \mathfrak{B}$ are bijective.*

As we mentioned earlier, we base our algorithm of reconstruction of the functions τ and q on the Krein accelerant method developed in [32].

Definition 1.5 *We say that a function $H \in L_2((-1, 1), M_r)$ is the accelerant if it is even (i.e., $H(-x) = H(x)$) and if for every $a \in [0, 1]$ the integral equation*

$$f(x) + \int_0^a H(x-t)f(t) dt = 0, \quad x \in (0, 1),$$

has no non-trivial solutions in the space L_2 . The set of all accelerants is denoted by $\mathfrak{H}_2$ and is endowed with the metric of the space $L_2((-1, 1), M_r)$. We shall write $\text{Re } \mathfrak{H}_2$ for the subset of $\mathfrak{H}_2$ of all Hermitian accelerants.

Spectral measure of the operator S_τ naturally generate Krein accelerants, as explained in the following theorem.

Theorem 1.6 *Take a sequence $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ satisfying condition (A_1) and set $\nu := \nu^{\mathbf{a}}$. Then the limit*

$$H_\nu(x) := \lim_{n \rightarrow \infty} \int_0^{\pi(n+1/2)} 2 \cos(2\lambda x) d(\nu - \nu_0)(\lambda), \quad x \in (-1, 1), \quad (1.10)$$

exists in the topology of the space $L_2((-1, 1), M_r)$. If, in addition, condition (A_3) holds, then the function H_ν belongs to $\text{Re } \mathfrak{H}_2$.

Conversely, every accelerant H determines a function in $\mathbf{L}_2$ in the following way. It is known (see Appendix A) that the associated Krein equation

$$R(x, t) + H(x - t) + \int_0^x R(x, \xi)H(\xi - t) d\xi = 0, \quad (x, t) \in \Omega, \quad (1.11)$$

has a unique solution R_H in the class $L_2(\Omega, M_r)$. We can now define a mapping Θ from $\mathfrak{H}_2$ to $\mathbf{L}_2$ given by the formula

$$[\Theta(H)](x) := H(x) + \int_0^x R_H(x, \xi)H(\xi) d\xi = -R_H(x, 0). \quad (1.12)$$

The functions $\Theta(H)$ and R_H are related to each other as follows:

Theorem 1.7 *Assume that $H \in \mathfrak{H}_2$, $R = R_H$, and $\tau = \Theta(H)$. Then*

$$\begin{aligned} K_{\tau, D}(x, t) &= \frac{1}{2} \left[R\left(x, \frac{x+t}{2}\right) - R\left(x, \frac{x-t}{2}\right) \right], \\ K_{\tau, N}(x, t) &= \frac{1}{2} \left[R\left(x, \frac{x+t}{2}\right) + R\left(x, \frac{x-t}{2}\right) \right], \end{aligned}$$

where $(x, t) \in \Omega$ and $K_{\tau, D}$ and $K_{\tau, N}$ are the kernels of (1.5).

Theorem 1.7 is an analogue of a theorem from the paper of Krein [32], where the accelerant theory was applied to the inverse scattering theory. Accelerants were used to solve the inverse spectral problems on finite intervals in the papers [33] and [34].

The following theorem describes some additional properties of the mapping Θ .

Theorem 1.8 *The mapping Θ is a homeomorphism between the metric spaces $\mathfrak{H}_2$ and $\mathbf{L}_2$. Moreover, if $H \in \mathfrak{H}_2$, then $H^* \in \mathfrak{H}_2$ and $\Theta(H^*) = [\Theta(H)]^*$.*

Finally, we show how the accelerants can be used to reconstruct τ and q from the corresponding spectral data.

Theorem 1.9 (i) *Assume that $\tau \in \text{Re } \mathbf{L}_2$ and that $\mathbf{a} := \mathbf{a}_\tau$ is the spectral data for the operator S_τ . Then $\tau = \Theta(H_\nu)$ for $\nu = \nu^\mathbf{a}$.*

(ii) *Assume that $q \in \text{Re } \mathfrak{M}$ and that $\mathbf{b} := \mathbf{b}_q$ is the spectral data for the operator T_q . Set $\nu := I\delta_0 + \mu^\mathbf{b}$ and $\tau := \Theta(H_\nu)$; then $q = b(\tau)$.*

According to Theorem 1.9, the reconstruction algorithm can proceed as follows. Given $\mathbf{a} \in \mathfrak{A}$, we construct a measure $\nu = \nu^\mathbf{a}$ via (1.9), which defines an accelerant $H = H_\nu$ by formula (1.10). Solving the Krein equation (1.11), we find the kernel R_H , which gives $\tau := \Theta(H)$ via (1.12). That τ so constructed is the matrix-function looked for follows from the fact that the matrix Sturm–Liouville operator S_τ has the spectral data $\mathbf{a}$ we have started with. As in [13], we visualize the reconstruction algorithm by means of the diagram

$$\mathbf{a} \xrightarrow[s_1]{(1.9)} \nu^\mathbf{a} = \nu \xrightarrow[s_2]{(1.10)} H_\nu = H \xrightarrow[s_3]{(1.11)} R_H \xrightarrow[s_4]{(1.12)} \Theta(H) = \tau.$$

In this diagram, s_j denotes the step number j . Steps s_1 , s_2 , and s_4 are trivial. The basic (and non-trivial) step is s_3 .

The procedure of reconstructing q looks similarly. Namely, given $\mathbf{b} \in \mathfrak{B}$, the corresponding potential $q \in \text{Re } \mathfrak{M}$ can be found following the steps in the next diagram:

$$\mathbf{b} \xrightarrow[s_1]{(1.9)} \mu^\mathbf{b} \xrightarrow[s_2]{\longrightarrow} I\delta_0 + \mu^\mathbf{b} = \nu \xrightarrow[s_3]{(1.10)} H_\nu = H \xrightarrow[s_4]{(1.11)} R_H \xrightarrow[s_5]{(1.12)} \Theta(H) = \tau \xrightarrow[s_6]{\longrightarrow} b(\tau) = q.$$

We observe that the spectral data for the operator T_q with $q = b(\tau)$ do not determine τ ; in fact, the Riccati equation $b(\tau) = q$ has many solutions, and by (1.2) each such solution generates the same operator T_q . Also, the assumption that $q \in \text{Re } \mathfrak{M}$ is not restrictive and was made only to simplify the presentation. Indeed, for every $q \in \text{Re } \mathbf{W}_2^{-1}$ the corresponding Dirichlet Sturm–Liouville operator T_q of (1.1) is bounded below and thus becomes positive after adding a suitable multiple cI of the unit matrix I ; it then follows that $q + cI$ belongs to $\text{Re } \mathfrak{M}$ (cf. [31]).

1.4. The structure of the paper

The paper is organised as follows. Section 2 deals with the direct spectral problem and consists of four parts. We establish basic properties of the operators T_q and S_τ (Theorem 2.4) in Subsection 2.1, find the asymptotics of eigenvalues and norming

constants (Theorem 2.5) in Subsection 2.2, study properties of the Weyl–Titchmarsh function in Subsection 2.3, and, finally, prove necessity of conditions of Theorems 1.1 and 1.2 in Subsection 2.4.

In Section 3 we study Krein accelerants. This section is divided in three parts. In the first two subsections we establish Theorems 1.7 and 1.8, and in Subsection 3.3 we prove Propositions 3.7 and 3.8 on operators $\mathcal{H}_o$ and $\mathcal{H}_e$, which play an important role in Section 4.

The inverse spectral problem is considered in Section 4, which consists of three parts. In Subsection 4.1 we study properties of the function H_ν (Theorem 1.6), in Subsection 4.2 we finish the proof of Theorems 1.1 and 1.2, and, finally, in Subsection 4.3 we establish Proposition 1.3 and Theorems 1.4 and 1.9.

There are two short appendices. Some information on the spaces used in the paper and well-known facts from the theory of factorization of Fredholm operators are gathered in Appendix A. Three auxiliary lemmata on orthogonal projectors are proved in Appendix B.

2. The direct spectral problem

2.1. Basic properties of the operators T_q and S_τ

In this subsection we prove self-adjointness of the operators T_q and S_τ in the case where $\tau \in \operatorname{Re} \mathbf{L}_2$ and $q \in \mathfrak{M}$ and also construct their resolvents and the resolutions of identity.

Recall that we have denoted by $\varphi(\cdot, \lambda, \tau)$ and $\psi(\cdot, \lambda, \tau)$ the matrix-valued solutions of the Cauchy problems (1.3) and (1.4). The results of [35] imply the following statement.

Theorem 2.1 *The problems (1.3) and (1.4) have unique solutions $\varphi(\cdot, \lambda, \tau)$ and $\psi(\cdot, \lambda, \tau)$ in the classes $\{u \in \mathbf{W}_2^1 \mid u_\tau^{[1]} \in \mathbf{W}_2^1\}$ and $\{u \in \mathbf{W}_2^1 \mid u_{-\tau}^{[1]} \in \mathbf{W}_2^1\}$, respectively. Moreover, the following statements hold:*

(i) *the functions $\varphi(\cdot, \lambda, \tau)$ and $\psi(\cdot, \lambda, \tau)$ are related to each other via*

$$\left(\frac{d}{dx} - \tau\right) \varphi(\cdot, \lambda, \tau) = \lambda \psi(\cdot, \lambda, \tau), \quad \left(\frac{d}{dx} + \tau\right) \psi(\cdot, \lambda, \tau) = -\lambda \varphi(\cdot, \lambda, \tau); \quad (2.1)$$

(ii) *for every $\tau \in \mathbf{L}_2$ there exist unique matrix-valued functions $K_{\tau,D}$ and $K_{\tau,N}$ belonging to the algebra G_2^+ (see Appendix A) such that for any $\lambda \in \mathbb{C}$ and $x \in [0, 1]$*

$$\begin{aligned} \varphi(x, \lambda, \tau) &= \sin \lambda x I + \int_0^x (\sin \lambda t) K_{\tau,D}(x, t) dt, \\ \psi(x, \lambda, \tau) &= \cos \lambda x I + \int_0^x (\cos \lambda t) K_{\tau,N}(x, t) dt; \end{aligned} \quad (2.2)$$

(iii) *the mappings $\mathbf{L}_2 \ni \tau \mapsto K_{\tau,D} \in G_2^+$ and $\mathbf{L}_2 \ni \tau \mapsto K_{\tau,N} \in G_2^+$ are continuous.*

Lemma 2.2 *Suppose that $\tau \in \mathbf{L}_2$ and $q = b(\tau)$. Then*

$$D(\mathbf{t}_{\tau,D}) = \{f \in W_{2,0}^1 \mid (-f'' + qf) \in L_2\}, \quad \mathbf{t}_{\tau,D}(f) = -f'' + qf,$$

where the derivative f'' and the product qf are interpreted in the distributional sense, i.e., as elements of the space W_2^{-1} . Moreover, if τ_1 is another element of $\mathbf{L}_2$, then

- (i) $\mathbf{t}_{\tau_1, D} = \mathbf{t}_{\tau, D} \iff b(\tau_1) = b(\tau)$;
- (ii) $K_{\tau_1, D} = K_{\tau, D} \iff b(\tau_1) = b(\tau)$;
- (iii) $K_{\tau_1, N} = K_{\tau, N} \iff \tau_1 = \tau$.

Proof. If $u \in \mathbf{W}_2^{-1}$ and $f \in W_{2,0}^1$, then we define the product $uf \in W_2^{-1}$ as

$$(uf)_j = \sum_{k=1}^r u_{jk} f_k, \quad f = (f_1, \dots, f_r), \quad u = (u_{jk})_{1 \leq j, k \leq r}.$$

Observe that for every $f \in W_{2,0}^1$ the equality $(\tau f)' = \tau' f + \tau f'$ holds in the distributional sense. If, therefore, $f \in D(\mathbf{t}_{\tau, D})$, then

$$\mathbf{t}_{\tau, D}(f) = -f'' + (\tau f)' - \tau f' + \tau^2 f = -f'' + qf \in L_2.$$

On the other hand, if $f \in W_{2,0}^1$ and $(-f'' + qf) \in L_2$, then taking into account the equality

$$-(f' - \tau f)' = (-f'' + qf) + \tau(f' - \tau f), \quad (2.3)$$

we obtain $(f' - \tau f) \in W_1^1$. Moreover, the function $f' - \tau f$ is bounded, and, using (2.3) again, we obtain that $(f' - \tau f) \in W_2^1$, i.e., that $f \in D(\mathbf{t}_{\tau, D})$. This establishes the claim about $\mathbf{t}_{\tau, D}$.

Part (i) is the straightforward consequence of the above considerations. Let us prove part (ii). Assume that $K_{\tau_1, D} = K_{\tau, D}$. Then $\varphi(\cdot, 0, \tau_1) = \varphi(\cdot, 0, \tau)$ in view of (2.2), and (2.1) yields the relations

$$(\tau_1 - \tau)\varphi(\cdot, \lambda, \tau) = \lambda[\psi(\cdot, \lambda, \tau) - \psi(\cdot, \lambda, \tau_1)], \quad (2.4)$$

$$\left(\frac{d}{dx} + \tau_1\right) \left(\frac{d}{dx} - \tau_1\right) \varphi(\cdot, \lambda, \tau) = \left(\frac{d}{dx} + \tau\right) \left(\frac{d}{dx} - \tau\right) \varphi(\cdot, \lambda, \tau). \quad (2.5)$$

Note that according to the Riemann–Lebesgue lemma for every $x_0 \in (0, 1]$ there exists $\lambda_0 \in \mathbb{R}$ such that $\det \varphi(x_0, \lambda_0, \tau) \neq 0$. Therefore we conclude from the equation (2.4) that the function $g = \tau_1 - \tau$ is absolutely continuous on the interval $(0, 1)$. Putting $\tau_1 = \tau + g$ in (2.5), we easily obtain the equality

$$(g' + g\tau + \tau g + g^2)\varphi(\cdot, \lambda, \tau) = 0, \quad \lambda \in \mathbb{C},$$

which implies that $g' + g\tau + \tau g + g^2 = 0$ and, thus, that $b(\tau_1) = b(\tau)$.

Now let $b(\tau_1) = b(\tau)$. Then the function $g = \tau_1 - \tau$ obeys $g' = \tau^2 - \tau_1^2 \in \mathbf{L}_1$ and thus is absolutely continuous on the interval $[0, 1]$. Next, the relation $g' + g\tau + \tau g + g^2 = 0$ yields equality (2.5). It follows from (2.5) and the definition of the functions $\varphi(\cdot, \lambda, \tau_1)$ and $\varphi(\cdot, \lambda, \tau)$ that they are solutions of the Cauchy problem

$$\left(\frac{d}{dx} + \tau_1\right) \left(\frac{d}{dx} - \tau_1\right) y = -\lambda^2 y, \quad y(0) = 0, \quad y_{\tau_1}^{[1]}(0) = \lambda I.$$

Thus $\varphi(\cdot, \lambda, \tau_1) = \varphi(\cdot, \lambda, \tau)$ for all $\lambda \in \mathbb{C}$. Therefore (see (2.2)),

$$\int_0^x (\sin \lambda t) [K_{\tau_1, D}(x, t) - K_{\tau, D}(x, t)] dt = 0, \quad x \in [0, 1], \quad \lambda \in \mathbb{C}.$$

Since λ and x are arbitrary, $K_{\tau_1, D} = K_{\tau, D}$.

It remains to prove (iii). If $\tau_1 = \tau$, then $K_{\tau_1, N} = K_{\tau, N}$ by the uniqueness claim of Theorem 2.1. Assume that $K_{\tau_1, N} = K_{\tau, N}$; then, according to (2.1) and (2.2),

$$\left(\frac{d}{dx} + \tau_1\right) \psi(x, 0, \tau_1) = \left(\frac{d}{dx} + \tau\right) \psi(x, 0, \tau) = 0, \quad \psi(\cdot, 0, \tau_1) = \psi(\cdot, 0, \tau),$$

and thus $(\tau_1 - \tau)\psi(\cdot, 0, \tau) = 0$. Therefore to prove part (iii) it is enough to show that the matrix $\psi(x, 0, \tau)$ is invertible for every $x \in [0, 1]$. Suppose the last statement is false. Then there exist $x_0 \in (0, 1]$ and $c \in \mathbb{C}^r \setminus \{0\}$ such that $\psi(x_0, 0, \tau)c = 0$. Therefore the function $f = \psi(\cdot, 0, \tau)c$ is a nonzero solution of the Cauchy problem $f' + \tau f = 0$, $f(x_0) = 0$, which is impossible. The proof is complete. $\square$

For every $\tau \in \mathbf{L}_2$, $\lambda \in \mathbb{C}$ and $x \in [0, 1]$ we put

$$\mathcal{W}(x, \lambda, \tau) = \begin{pmatrix} \psi(x, \lambda, \tau) & \varphi(x, \lambda, -\tau) \\ -\varphi(x, \lambda, \tau) & \psi(x, \lambda, -\tau) \end{pmatrix}, \quad \mathcal{Q}(x, \lambda, \tau) = \begin{pmatrix} -\tau(x) & \lambda I \\ -\lambda I & \tau(x) \end{pmatrix}.$$

It follows from (2.1) that $\frac{d}{dx}\mathcal{W}(x, \lambda, \tau) = \mathcal{Q}(x, \lambda, \tau)\mathcal{W}(x, \lambda, \tau)$, so that

$$\frac{d}{dx}\mathcal{W}^*(x, \bar{\lambda}, -\tau^*)\mathcal{W}(x, \lambda, \tau) = 0.$$

Therefore the matrices $\mathcal{W}^*(x, \bar{\lambda}, -\tau^*)$ and $\mathcal{W}(x, \lambda, \tau)$ are inverse to each other, which yields the relations

$$\begin{aligned} \varphi(x, \lambda, \tau)\varphi^*(x, \bar{\lambda}, -\tau^*) + \psi(x, \lambda, -\tau)\psi^*(x, \bar{\lambda}, \tau^*) &\equiv I, \\ \varphi(x, \lambda, \tau)\psi^*(x, \bar{\lambda}, -\tau^*) - \psi(x, \lambda, -\tau)\varphi^*(x, \bar{\lambda}, \tau^*) &\equiv 0. \end{aligned} \tag{2.6}$$

Let us denote by $\Phi_\tau(\lambda)$ and $\Psi_\tau(\lambda)$ ($\lambda \in \mathbb{C}$) the operators acting from $\mathbb{C}^r$ to L_2 by the formulas

$$[\Phi_\tau(\lambda)c](x) := \sqrt{2}\varphi(x, \lambda, \tau)c, \quad [\Psi_\tau(\lambda)c](x) := \sqrt{2}\psi(x, \lambda, \tau)c. \tag{2.7}$$

Taking into consideration (2.2), we obtain, for $\lambda \in \mathbb{C}$,

$$\Phi_\tau(\lambda) = (\mathcal{I} + \mathcal{K}_{\tau, D})\Phi_0(\lambda), \quad \Psi_\tau(\lambda) = (\mathcal{I} + \mathcal{K}_{\tau, N})\Psi_0(\lambda), \tag{2.8}$$

where $\mathcal{K}_{\tau, D}$ and $\mathcal{K}_{\tau, N}$ are integral operators with kernels $K_{\tau, D}$ and $K_{\tau, N}$ respectively and $\mathcal{I}$ is the identity operator in the algebra $\mathcal{B}(L_2)$, which is an algebra of bounded linear operators acting in L_2 . Note that since $K_{\tau, D}$ and $K_{\tau, N}$ belong to G_2^+ , the operators $\mathcal{K}_{\tau, D}$ and $\mathcal{K}_{\tau, N}$ belong to the algebra $\mathfrak{G}_2^+$ (see Appendix A), and hence they are Volterra operators [36, Ch. IV].

Lemma 2.3 *Let $\tau \in \mathbf{L}_2$. Then the following statements hold:*

(i) *the operator functions $\lambda \mapsto \Phi_\tau(\lambda)/\lambda$ and $\lambda \mapsto \Psi_\tau(\lambda)$ are analytic in $\mathbb{C}$; moreover,*

$$\begin{aligned} \ker \Phi_\tau(\lambda) &= \{0\}, & \text{Ran } \Phi_\tau^*(\lambda) &= \mathbb{C}^r, & \lambda &\in \mathbb{C} \setminus \{0\}, \\ \ker \Psi_\tau(\lambda) &= \{0\}, & \text{Ran } \Psi_\tau^*(\lambda) &= \mathbb{C}^r, & \lambda &\in \mathbb{C}, \end{aligned} \tag{2.9}$$

$$\ker(T_q - \lambda^2 \mathcal{I}) = \Phi_\tau(\lambda)\mathcal{E}_\lambda, \quad \ker(S_\tau - \lambda^2 \mathcal{I}) = \Psi_\tau(\lambda)\mathcal{E}_\lambda, \quad \lambda \in \mathbb{C}, \tag{2.10}$$

where $\mathcal{E}_\lambda := \ker \varphi(1, \lambda, \tau)$;

(ii) the operator functions $\lambda \mapsto \varphi(1, \lambda, \tau)^{-1}$ and

$$\lambda \mapsto m_\tau(\lambda) = -\varphi(1, \lambda, \tau)^{-1}\psi(1, \lambda, -\tau) \quad (2.11)$$

are meromorphic in $\mathbb{C}$. Moreover, $m_0(\lambda) = -\cot \lambda I$ and

$$\|m_\tau(\lambda) + \cot \lambda I\| = o(1) \quad (2.12)$$

as $\lambda \rightarrow \infty$ within the domain $\mathcal{O} = \{z \in \mathbb{C} \mid \forall n \in \mathbb{Z} \quad |z - \pi n| > 1\}$.

Proof. Part (i) obviously follows from (2.7) and (2.8). From (2.2) we obtain that

$$\begin{aligned} \varphi(1, \lambda, \tau) &= \sin \lambda I + \int_0^1 (\sin \lambda t) K_{\tau,D}(1, t) dt, \\ \psi(1, \lambda, -\tau) &= \cos \lambda I + \int_0^1 (\cos \lambda t) K_{-\tau,N}(1, t) dt. \end{aligned} \quad (2.13)$$

Since $K_{\tau,D}, K_{-\tau,N} \in G_2^+$, the functions $K_{\tau,D}(1, \cdot)$ and $K_{-\tau,N}(1, \cdot)$ belong to $\mathbf{L}_2$. The relations

$$\lim_{|\lambda| \rightarrow \infty} e^{-|Im \lambda|} \int_0^1 f(x) \cos \lambda x dx = \lim_{|\lambda| \rightarrow \infty} e^{-|Im \lambda|} \int_0^1 f(x) \sin \lambda x dx = 0$$

for $f \in L_1(0, 1)$ that refine the classical Riemann–Lebesgue theorem [37, Ch.1] are proved in [1, Ch.1]; thus

$$\lim_{|\lambda| \rightarrow \infty} e^{-|Im \lambda|} \|\varphi(1, \lambda, \tau) - \sin \lambda I\| = \lim_{|\lambda| \rightarrow \infty} e^{-|Im \lambda|} \|\psi(1, \lambda, -\tau) - \cos \lambda I\| = 0.$$

In particular, $\varphi(1, \lambda, \tau)$ is invertible for all $\lambda \in \mathcal{O}$ large enough, so that m_τ is meromorphic and the relation (2.12) holds. $\square$

Theorem 2.4 *Let $\tau \in Re \mathbf{L}_2$ and $q = b(\tau)$. Then the following statements hold:*

- (i) *The operators S_τ and T_q are self-adjoint; moreover, $S_\tau \geq 0$ and $T_q > 0$.*
- (ii) *The spectra $\sigma(S_\tau)$ and $\sigma(T_q)$ consist of isolated eigenvalues and*

$$\sigma(S_\tau) = \{\lambda^2 \mid \ker \varphi(1, \lambda, \tau) \neq \{0\}\} = \sigma(T_q) \cup \{0\}. \quad (2.14)$$

- (iii) *Let $\lambda_j = \lambda_j(\tau)$ and $P_{j,\tau}$ (resp. $Q_{j,q}$) be the orthogonal projector on the eigensubspace $\ker(S_\tau - \lambda_j^2 \mathcal{I})$ (resp. $\ker(T_q - \lambda_j^2 \mathcal{I})$); then*

$$\sum_{j=0}^{\infty} P_{j,\tau} = \mathcal{I}, \quad \sum_{k=1}^{\infty} Q_{k,q} = \mathcal{I}. \quad (2.15)$$

- (iv) *The norming constants $\alpha_j = \alpha_j(\tau)$ (see Introduction) satisfy the relations $\alpha_0 > 0$ and $\alpha_j \geq 0$, $j \in \mathbb{N}$. Moreover, for $j \in \mathbb{Z}_+$ and $k \in \mathbb{N}$ we have*

$$P_{j,\tau} = \Psi_\tau(\lambda_j) \alpha_j \Psi_\tau^*(\lambda_j), \quad Q_{k,q} = \Phi_\tau(\lambda_k) \alpha_k \Phi_\tau^*(\lambda_k), \quad (2.16)$$

where $\Phi_\tau^*(\lambda) = [\Phi_\tau(\lambda)]^*$, $\Psi_\tau^*(\lambda) = [\Psi_\tau(\lambda)]^*$ (see (2.7)).

- (v) *If $\tau_1 \in Re \mathbf{L}_2$ and $b(\tau_1) = b(\tau)$, then $\lambda_j(\tau_1) = \lambda_j(\tau)$ and $\alpha_j(\tau_1) = \alpha_j(\tau)$ for every $j \in \mathbb{N}$.*

Proof. Writing $S = S_\tau$ and $T = T_q$ for short and integrating by parts, we get

$$(Sf|g) = (f' - \tau f|g' - \tau g) = (f|Sg), \quad f, g \in D(S),$$

$$(Tf|g) = (f' - \tau f|g' - \tau g) = (f|Tg), \quad f, g \in D(T),$$

where $(\cdot|\cdot)$ is the scalar product in the space L_2 . Therefore the operators S and T are symmetric and nonnegative. Suppose that $(Tf|f) = 0$ for some $f \in D(T)$. Then f is a solution of the Cauchy problem $f' - \tau f = 0$, $f(0) = 0$. The uniqueness theorem then gives $f = 0$, and, therefore $T > 0$.

Take now an arbitrary $f \in L_2$ and a point $\lambda \in \mathbb{C} \setminus \{0\}$ for which the matrix $\varphi(1, \lambda, \tau)$ is nonsingular. The function

$$\begin{aligned} g(x) = [X(\lambda)f](x) &:= \frac{\varphi(x, \lambda, \tau)}{\lambda} \int_x^1 \psi^*(t, \bar{\lambda}, -\tau) f(t) dt \\ &+ \frac{\psi(x, \lambda, -\tau)}{\lambda} \int_0^x \varphi^*(t, \bar{\lambda}, \tau) f(t) dt \end{aligned}$$

vanishes at $x = 0$, belongs to the domain of the differential expression $\mathbf{t}_\tau$, and solves the equation $\mathbf{t}_\tau g = \lambda^2 g + f$, as can be verified directly by using the relations (2.1) and (2.6). A generic solution of the above differential equation that vanishes at $x = 0$ takes therefore the form $\varphi(x, \lambda, \tau)c + g$ for some vector $c \in \mathbb{C}^r$; the choice

$$c := \frac{m_\tau(\lambda)}{\lambda} \int_0^1 \varphi^*(t, \bar{\lambda}, \tau) f(t) dt$$

makes this solution to vanish at the point $x = 1$ as well. This implies that the point λ^2 is a resolvent point of the operator T and that the resolvent of T is given by

$$(T - \lambda^2 \mathcal{I})^{-1} = \frac{1}{2\lambda} \Phi_\tau(\lambda) m_\tau(\lambda) \Phi_\tau^*(\bar{\lambda}) + X(\lambda).$$

Similar arguments show that the resolvent of the operator S at the point λ^2 equals

$$(S - \lambda^2 \mathcal{I})^{-1} = \frac{1}{2\lambda} \Psi_\tau(\lambda) m_\tau(\lambda) \Psi_\tau^*(\bar{\lambda}) - Y(\lambda),$$

where the operator Y is given via

$$[Y(\lambda)f](x) = \frac{\varphi(x, \lambda, -\tau)}{\lambda} \int_0^x \psi^*(t, \bar{\lambda}, \tau) f(t) dt + \frac{\psi(x, \lambda, \tau)}{\lambda} \int_x^1 \varphi^*(t, \bar{\lambda}, -\tau) f(t) dt.$$

The above formulas show that S and T have compact resolvents and thus (i)–(iii) follow.

Recall that $-\alpha_j(\tau)$ is the residue of the Weyl–Titchmarsh function m_τ at the point $\lambda_j(\tau)$, $j \in \mathbb{N}$. Taking $\varepsilon > 0$ small enough, we get

$$\begin{aligned} P_{j,\tau} &= \frac{-1}{2\pi i} \oint_{|\zeta - \lambda_j^2| = \varepsilon} (S - \zeta \mathcal{I})^{-1} d\zeta = \frac{-1}{2\pi i} \oint_{|\lambda - \lambda_j| = \varepsilon} 2\lambda (S - \lambda^2 \mathcal{I})^{-1} d\lambda \\ &= \frac{-1}{2\pi i} \oint_{|\lambda - \lambda_j| = \varepsilon} \Psi_\tau(\lambda) m_\tau(\lambda) \Psi_\tau^*(\lambda) d\lambda = \Psi_\tau(\lambda_j) \alpha_j \Psi_\tau^*(\lambda_j) \end{aligned}$$

for every $j \in \mathbb{N}$. Similarly, we obtain that

$$P_{0,\tau} = \Psi_\tau(0) \alpha_0 \Psi_\tau^*(0), \quad Q_{j,q} = \Phi_\tau(\lambda_j) \alpha_j \Phi_\tau^*(\lambda_j), \quad j \in \mathbb{N}.$$

Recalling (2.9), we see that $\alpha_j \geq 0$ for all $j \in \mathbb{Z}_+$. Let us prove that $\alpha_0 > 0$. Assume the contrary; then

$$\ker S = \operatorname{Ran} P_{0,\tau} = \operatorname{Ran} [\Psi_\tau(0)\alpha_0\Psi_\tau^*(0)] \neq \operatorname{Ran} \Psi_\tau(0).$$

On the other hand, (2.10) on account of the equality $\ker \varphi(1, 0, \tau) = \mathbb{C}^r$ yields $\ker S = \operatorname{Ran} \Psi_\tau(0)$. The contradiction derived shows that $\alpha_0 > 0$.

It remains to prove (v). Let $\tau_1 \in \operatorname{Re} \mathbf{L}_2$ be such that $b(\tau_1) = b(\tau)$. It follows from Lemma 2.2 that $K_{\tau_1,D} = K_{\tau,D}$, and thus $\varphi(\cdot, \cdot, \tau_1) = \varphi(\cdot, \cdot, \tau)$. Therefore, in view of (2.14) we get $\lambda_j(\tau_1) = \lambda_j(\tau)$ for all $j \in \mathbb{N}$ and, moreover, $\Phi_{\tau_1}(\cdot) = \Phi_\tau(\cdot)$. It follows now from (2.16) that

$$\Phi_\tau(\lambda_j)\alpha_j(\tau_1)\Phi_\tau^*(\lambda_j) = Q_{j,q} = \Phi_\tau(\lambda_j)\alpha_j(\tau)\Phi_\tau^*(\lambda_j), \quad j \in \mathbb{N}.$$

Hence, using (2.9), we obtain that $\alpha_j(\tau_1) = \alpha_j(\tau)$ for all $j \in \mathbb{N}$. The proof is complete. $\square$

2.2. The asymptotics of eigenvalues and norming constants

The main result of this section is following theorem.

Theorem 2.5 *Let $\tau \in \operatorname{Re} \mathbf{L}_2$. Then for the sequence $\mathbf{a} = \mathbf{a}_\tau$ the condition (A_1) holds.*

First we prove two lemmas. In the sequel, we shall use the following notation. If $(\lambda_j)_{j \in \mathbb{Z}_+}$ is a strictly increasing sequence of nonnegative numbers and $(\alpha_j)_{j \in \mathbb{Z}_+}$ is a sequence in M_r^+ , then

$$\beta_n := I - \sum_{\lambda_k \in \Delta_n} \alpha_k, \quad \tilde{\lambda}_j := \lambda_j - \pi n, \quad \lambda_j \in \Delta_n, \quad n \in \mathbb{N}, \quad (2.17)$$

with Δ_n defined in Subsection 1.3.

Lemma 2.6 *Let $\tau \in \operatorname{Re} \mathbf{L}_2$ and $\lambda_j = \lambda_j(\tau)$, $j \in \mathbb{Z}_+$. Then*

$$\sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} |\tilde{\lambda}_j|^2 < \infty, \quad \sup_{n \in \mathbb{N}} \sum_{\lambda_j \in \Delta_n} 1 < \infty. \quad (2.18)$$

Proof. Let us fix $\tau \in \operatorname{Re} \mathbf{L}_2$ and note that the numbers $\lambda_j(\tau)$, $j \in \mathbb{Z}_+$, are nonnegative zeros of an entire function $g(\lambda) := \det \varphi(1, \lambda, \tau)$, $\lambda \in \mathbb{C}$. In view of (2.13) the function g is odd. Taking into account (2.14), we conclude that zeros of the function g are all real. The function $\lambda \mapsto \varphi(1, \lambda, \tau)$ belongs to the following class of function $\mathbb{C} \rightarrow M_r$:

$$\mathcal{F}_f(\lambda) := \sin \lambda I + \int_{-1}^1 f(t) e^{i\lambda t} dt, \quad \lambda \in \mathbb{C},$$

where $f \in L_2((-1, 1), M_r)$. Indeed, it follows from (2.13) that $\varphi(1, \cdot, \tau) = \mathcal{F}_f$ for $f(t) = (\operatorname{sign} t) K_{\tau,D}(1, |t|)/(2i)$. It is shown in the paper [38] that the set of zeros of a function $\det \mathcal{F}_f$ with $\mathcal{F}_f$ as above can be indexed (counting multiplicities) by the set $\mathbb{Z}$ so that the corresponding sequence $(\omega_n)_{n \in \mathbb{Z}}$ has the asymptotics

$$\omega_{kr+j} = \pi k + \hat{\omega}_{j,k}, \quad k \in \mathbb{Z}, \quad j = 0, \dots, r-1,$$

where the sequences $(\widehat{\omega}_{j,k})_{k \in \mathbb{Z}}$ belong to $\ell_2(\mathbb{Z})$. Therefore, (2.18) follows, and the proof is complete. $\square$

Lemma 2.7 *Let the operators $A(z)$ and $B(z)$ ($z \in \mathbb{C}$) act from $\mathbf{L}_2$ to M_r by the formulas*

$$A(z)f = \sqrt{2} \int_0^1 (\sin zt) f(t) dt, \quad B(z)g = \sqrt{2} \int_0^1 (\cos zt) g(t) dt. \quad (2.19)$$

Then for every $f_1, f_2 \in \mathbf{L}_2$ and $\lambda \in \mathbb{D} := \{\lambda \in \mathbb{C} \mid |\lambda| \leq 1\}$ we have

$$\sum_{n=1}^{\infty} (\|A(\pi n + \lambda)f_1\|^2 + \|B(\pi n + \lambda)f_2\|^2) \leq 16r^2(\|f_1\|_{\mathbf{L}_2}^2 + \|f_2\|_{\mathbf{L}_2}^2). \quad (2.20)$$

Proof. Since $\{\sqrt{2} \sin \pi nt\}_{n \in \mathbb{N}}$ and $\{\sqrt{2} \cos \pi nt\}_{n \in \mathbb{N}}$ form orthonormal systems of $L_2(0, 1)$, it follows that

$$\sum_{n=1}^{\infty} (\|A(\pi n)f_1\|^2 + \|B(\pi n)f_2\|^2) \leq 2r^2(\|f_1\|_{\mathbf{L}_2}^2 + \|f_2\|_{\mathbf{L}_2}^2). \quad (2.21)$$

Relations (2.19) yield

$$A(\pi n + \lambda)f_1 = B(\pi n)f_{11} + A(\pi n)f_{12}, \quad B(\pi n + \lambda)f_2 = B(\pi n)f_{22} - A(\pi n)f_{21}, \quad (2.22)$$

where

$$f_{j1}(t) = (\sin \lambda t)f_j(t), \quad f_{j2}(t) = (\cos \lambda t)f_j(t), \quad t \in (0, 1), \quad j = 1, 2.$$

Since $|\sin \lambda|^2 + |\cos \lambda|^2 \leq (\sinh 1 + \cosh 1)^2 \leq 8$ for $\lambda \in \mathbb{D}$, we have

$$\|f_{j1}\|_{\mathbf{L}_2}^2 + \|f_{j2}\|_{\mathbf{L}_2}^2 \leq 8\|f_j\|_{\mathbf{L}_2}^2, \quad j = 1, 2.$$

Hence, using (2.22) and (2.21), we arrive at (2.20). $\square$

Proof of Theorem 2.5 Let $\tau \in \operatorname{Re} \mathbf{L}_2$ and $\lambda_j = \lambda_j(\tau), \alpha_j = \alpha_j(\tau)$. According to Lemma 2.6, it remains to show that $\sum_{n=1}^{\infty} \|\beta_n\|^2 < \infty$. Set

$$\mathbb{T}_n := \{z \in \mathbb{C} \mid |\lambda - \pi n| = 1\}, \quad \mathbb{D}_n := \{z \in \mathbb{C} \mid |\lambda - \pi n| \leq 1\}, \quad n \in \mathbb{Z}_+,$$

$$f_1 := 2^{-1/2}K_{\tau,D}(1, \cdot), \quad f_2 := 2^{-1/2}K_{-\tau,N}(1, \cdot).$$

Since $K_{\tau,D}, K_{-\tau,N} \in G_2^+$, we have $f_1, f_2 \in \mathbf{L}_2$. It follows from (2.13) and (2.19) that

$$\varphi(1, \lambda, \tau) = \sin \lambda I + A(\lambda)f_1, \quad \psi(1, \lambda, -\tau) = \cos \lambda I + B(\lambda)f_2.$$

Hence, using (2.11), we obtain the equality

$$m_\tau(\lambda) + \cot(\lambda)I = \varphi(1, \lambda, \tau)^{-1}[(\cot \lambda)A(\lambda)f_1 - B(\lambda)f_2]. \quad (2.23)$$

In view of (2.18) and (2.19), we can choose $n_0 \in \mathbb{N}$, such that

$$\sum_{n=n_0}^{\infty} \sum_{\lambda_j \in \Delta_n} |\tilde{\lambda}_j|^2 < 1, \quad \sup_{n \geq n_0} \sup_{\lambda \in \mathbb{D}_n} \|A(\lambda)f_1\| \leq \frac{1}{4}. \quad (2.24)$$

Note that for every $\lambda \in \mathbb{T}_0$

$$|\sin \lambda| \geq 1 - \sum_{n=1}^{\infty} \frac{1}{(2n+1)!!} \geq 1 - \sum_{n=1}^{\infty} \frac{1}{3^n} = \frac{1}{2},$$

whence $|\sin \lambda| \geq \frac{1}{2}$ and $|\cot \lambda| \leq \sqrt{3}$ for $\lambda \in \mathbb{T}_n$, $n \in \mathbb{Z}_+$. Therefore estimates (2.24) yield the following inequalities for $\lambda \in \mathbb{T}_n$ with $n \geq n_0$:

$$\|\varphi(1, \lambda, \tau)^{-1}\| \leq |\sin \lambda|^{-1} (1 - |\sin \lambda|^{-1} \|A(\lambda) f_1\|)^{-1} \leq 4. \quad (2.25)$$

For $n \geq n_0$ the function m_τ has no poles on the circle $\mathbb{T}_n$ and $\{\lambda_j \mid \lambda_j \in \Delta_n\} \subset \mathbb{D}_n$. Hence (1.6) implies that

$$\beta_n = \frac{1}{2\pi i} \oint_{\mathbb{T}_n} (m_\tau(\lambda) + \cot \lambda I) d\lambda, \quad n \geq n_0. \quad (2.26)$$

Using (2.23) and (2.25), we get

$$\|m_\tau(\lambda) + \cot \lambda I\|^2 \leq 96(\|A(\lambda) f_1\|^2 + \|B(\lambda) f_2\|^2), \quad \lambda \in \mathbb{T}_n, \quad n \geq n_0.$$

It follows now from (2.26) that

$$\|\beta_n\|^2 \leq \frac{96}{2\pi} \int_{\mathbb{T}_0} \{\|A(\pi n + e^{it}) f_1\|^2 + \|B(\pi n + e^{it}) f_2\|^2\} dt, \quad n \geq n_0,$$

and in view of (2.20) we get $\sum_{n=1}^{\infty} \|\beta_n\|^2 < \infty$. $\square$

2.3. The Weyl–Titchmarsh function

Proposition 2.8 *Let $\tau \in \text{Re } \mathbf{L}_2$. Then m_τ is a Herglotz function and*

$$m_\tau(\lambda) = 2\lambda \int_0^\infty \frac{d\nu_\tau(\xi)}{\xi^2 - \lambda^2}, \quad \lambda \in \mathbb{C}. \quad (2.27)$$

Proof. That m_τ is Herglotz is clear, so we only need to prove (2.27). Set

$$h(\lambda) := m_\tau(\lambda) - 2\lambda \int_0^\infty \frac{d\nu_\tau(\xi)}{\xi^2 - \lambda^2}, \quad \lambda \in \mathbb{C}.$$

It follows from (1.6) and (1.7) that h is entire. To show that $h = 0$ it suffices to justify the estimate $\|h(\lambda)\| = o(1)$ as λ goes to infinity within the domain

$$\mathcal{O} = \{z \in \mathbb{C} \mid \forall n \in \mathbb{Z} \quad |z - \pi n| > 1\}.$$

We set

$$g(\lambda) := \cot \lambda I + 2\lambda \int_0^\infty \frac{d\nu_\tau(\xi)}{\xi^2 - \lambda^2}, \quad \lambda \in \mathbb{C}.$$

In virtue of (2.12), it suffices to prove the estimate

$$\|g(\lambda)\| = o(1), \quad \mathcal{O} \ni \lambda \rightarrow \infty. \quad (2.28)$$

Since

$$\cot \lambda = \frac{1}{\lambda} + \sum_{n=1}^{\infty} \frac{2\lambda}{\lambda^2 - (\pi n)^2}, \quad \lambda \in \mathbb{C},$$

we find that

$$g(\lambda) = \frac{1}{\lambda}(I - 2\alpha_0) + \sum_{n=1}^{\infty} \frac{2\lambda I}{\lambda^2 - (\pi n)^2} - \sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} \frac{2\lambda \alpha_j}{\lambda^2 - \lambda_j^2}.$$

Let

$$g_n(\lambda) = \frac{2\lambda I}{\lambda^2 - (\pi n)^2} - \sum_{\lambda_j \in \Delta_n} \frac{2\lambda \alpha_j}{\lambda^2 - \lambda_j^2}, \quad n \in \mathbb{N}.$$

It is easy to verify that for arbitrary $\lambda_j \in \Delta_n$ such that $|\tilde{\lambda}_j| \leq 1/2$,

$$\left| \frac{2\lambda}{\lambda^2 - (\pi n)^2} - \frac{2\lambda}{\lambda^2 - \lambda_j^2} \right| \leq \frac{2|\tilde{\lambda}_j|}{|\lambda - \pi n|} + \frac{2|\tilde{\lambda}_j|}{|\lambda + \pi n|}, \quad \lambda \in \mathcal{O}.$$

Therefore, for big enough n and $\lambda \in \mathcal{O}$,

$$\|g_n(\lambda)\| \leq \left(\|\beta_n\| + \sum_{\lambda_j \in \Delta_n} 2|\tilde{\lambda}_j| \|\alpha_j\| \right) \left(\frac{1}{|\lambda - \pi n|} + \frac{1}{|\lambda + \pi n|} \right).$$

Taking into account Theorem 2.5 and using the Cauchy–Bunyakovski inequality, we get that there exists a constant C such that, for all big enough k and $\lambda \in \mathcal{O}$,

$$\sum_{n=k}^{\infty} \|g_n(\lambda)\| \leq C \left(\sum_{n=k}^{\infty} \|\beta_n\|^2 + \sum_{n=k}^{\infty} \sum_{\lambda_j \in \Delta_n} |\tilde{\lambda}_j|^2 \right)^{1/2} \left(\sum_{n \in \mathbb{Z}} \frac{1}{|\lambda - \pi n|^2} \right)^{1/2}. \quad (2.29)$$

Since

$$\sum_{n \in \mathbb{Z}} \frac{1}{|\lambda - \pi n|^2} \leq 2 + \frac{2}{\pi^2} \sum_{n=1}^{\infty} \frac{1}{n^2} \leq 3, \quad \lambda \in \mathcal{O},$$

and $\|g_n(\lambda)\| = O(1/\lambda)$ as $\lambda \rightarrow \infty$, (2.28) follows from (2.29). The proof is complete. $\square$

2.4. Necessity parts of Theorems 1.1 and 1.2

The theorem stated below implies necessity in Theorems 1.1 and 1.2.

Theorem 2.9 *Let $\tau \in \text{Re } \mathbf{L}_2$. Then for a sequence $\mathbf{a}_\tau$ the conditions (A_1) – (A_4) hold.*

First we prove four auxiliary but nonetheless important lemmas.

Lemma 2.10 *Assume that $\tau \in \mathbf{L}_2$ and let $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ satisfy condition (A_1) . Then*

$$\sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} \|\Psi_\tau(\lambda_j) - \Psi_0(\pi n)\|^2 < \infty, \quad \sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} \|\Phi_\tau(\lambda_j) - \Phi_0(\pi n)\|^2 < \infty. \quad (2.30)$$

Proof. We prove the first inequality; the second one can be proved similarly. Recalling the definitions in (2.7)–(2.8), we conclude that for every λ and ξ in $\mathbb{R}$ one has

$$\|\Psi_\tau(\lambda)\| \leq C, \quad \|\Psi_\tau(\lambda) - \Psi_\tau(\xi)\| \leq C|\lambda - \xi|$$

with $C = \sqrt{2}(1 + \|\mathcal{K}_{\tau,N}\|)$. By virtue of (2.17) we derive the estimate

$$\sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} \|\Psi_{\tau}(\lambda_j) - \Psi_{\tau}(\pi n)\|^2 \leq C^2 \sum_{n=1}^{\infty} |\tilde{\lambda}_j|^2 < \infty. \quad (2.31)$$

We note that the operator $\mathcal{K}_{\tau,N}$ belongs to the Hilbert–Schmidt class $\mathcal{B}_2$ and that the sequence $(P_{n,0})_{n \in \mathbb{Z}_+}$ consists of pairwise orthogonal projectors. Therefore,

$$\sum_{n=1}^{\infty} \|\mathcal{K}_{\tau,N} P_{n,0}\|^2 \leq \sum_{n=1}^{\infty} \|\mathcal{K}_{\tau,N} P_{n,0}\|_{\mathcal{B}_2}^2 \leq \|\mathcal{K}_{\tau,N}\|_{\mathcal{B}_2}^2.$$

It follows from (2.8) that $\Psi_{\tau}(\pi n) - \Psi_0(\pi n) = \mathcal{K}_{\tau,N} \Psi_0(\pi n)$. Since $\|\Psi_0(\pi n)\| = 1$ and $P_{n,0} \Psi_0(\pi n) = \Psi_0(\pi n)$,

$$\sum_{n=1}^{\infty} \|\Psi_{\tau}(\pi n) - \Psi_0(\pi n)\|^2 \leq \sum_{n=1}^{\infty} \|\mathcal{K}_{\tau,N} P_{n,0} \Psi_0(\pi n)\|^2 \leq \|\mathcal{K}_{\tau,N}\|_{\mathcal{B}_2}^2. \quad (2.32)$$

Combining (2.31) and (2.32) with the uniform bound $\sup_{n \in \mathbb{N}} \sum_{\lambda_j \in \Delta_n} 1 < \infty$ satisfied due to (A_1) , we obtain the inequality (2.30) for the operators Ψ_{τ} . $\square$

Lemma 2.11 *Assume that $\tau \in \mathbf{L}_2$ and let $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ satisfy condition (A_1) . Then for every $f \in L_2$ we get*

$$\sum_{j=0}^{\infty} \|\alpha_j \Psi_{\tau}^*(\lambda_j) f\|_{\mathbb{C}^r}^2 < \infty, \quad \sum_{j=1}^{\infty} \|\alpha_j \Phi_{\tau}^*(\lambda_j) f\|_{\mathbb{C}^r}^2 < \infty. \quad (2.33)$$

Furthermore, for every sequence $(c_j) \in \ell_2(\mathbb{Z}_+, \mathbb{C}^r)$ the series $\sum_{j=0}^{\infty} \Psi_{\tau}(\lambda_j) c_j$ and $\sum_{j=1}^{\infty} \Phi_{\tau}(\lambda_j) c_j$ converge in the space L_2 .

Proof. We shall only prove the first inequality in (2.33) and convergence of the series $\sum_{j=0}^{\infty} \Psi_{\tau}(\lambda_j) c_j$; the other statements are justified analogously. Since the norms of the matrices α_j are uniformly bounded, it suffices to prove the inequality $\sum_{j=0}^{\infty} \|\Psi_{\tau}^*(\lambda_j) f\|_{\mathbb{C}^r}^2 < \infty$. Observe that

$$\begin{aligned} \sum_{j=0}^{\infty} \|\Psi_{\tau}^*(\lambda_j) f\|_{\mathbb{C}^r}^2 &= \sum_{n=0}^{\infty} \sum_{\lambda_j \in \Delta_n} \|\Psi_{\tau}^*(\lambda_j) f\|_{\mathbb{C}^r}^2 \\ &\leq 2 \sum_{n=0}^{\infty} \sum_{\lambda_j \in \Delta_n} \left\{ \|[\Psi_{\tau}^*(\lambda_j) - \Psi_0^*(\pi n)] f\|_{\mathbb{C}^r}^2 + \|\Psi_0^*(\pi n) f\|_{\mathbb{C}^r}^2 \right\} \end{aligned}$$

and that

$$\sum_{n=1}^{\infty} \|\Psi_0^*(\pi n) f\|_{\mathbb{C}^r}^2 \leq \|f\|_{\mathbb{C}^r}^2$$

by the Bessel inequality for the orthonormal system $\{\sqrt{2} \cos \pi n x\}_{n \in \mathbb{N}}$. The desired inequality follows now from Lemma 2.10 and the fact that, by virtue of (A_1) , $\sup_{n \in \mathbb{N}} \sum_{\lambda_j \in \Delta_n} 1 < \infty$.

To prove convergence of the series $\sum_{j=0}^{\infty} \Psi_{\tau}(\lambda_j) c_j$, we similarly write

$$\sum_{j=0}^{\infty} \Psi_{\tau}(\lambda_j) c_j = \sum_{n=0}^{\infty} \sum_{\lambda_j \in \Delta_n} [\Psi_{\tau}(\lambda_j) - \Psi_0(\pi n)] c_j + \sum_{n=0}^{\infty} \sum_{\lambda_j \in \Delta_n} \Psi_0(\pi n) c_j.$$

Since $(c_j) \in \ell_2(\mathbb{Z}_+, \mathbb{C}^r)$, the first series converges in L_2 due to (2.30) and the Cauchy–Bunyakowski inequality, while the second one due to the fact that the system $\{\sqrt{2} \cos \pi n x\}_{n \in \mathbb{N}}$ is orthonormal in $L_2(0, 1)$. $\square$

Lemma 2.12 *Assume that $\tau \in \mathbf{L}_2$, let $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ satisfy condition (A_1) , and set*

$$\widehat{P}_j = \Psi_{\tau}(\lambda_j) \alpha_j \Psi_{\tau}^*(\lambda_j), \quad \widehat{Q}_j = \Phi_{\tau}(\lambda_j) \alpha_j \Phi_{\tau}^*(\lambda_j), \quad j \in \mathbb{N}.$$

Then the series $\sum_{j=1}^{\infty} \widehat{P}_j$ and $\sum_{j=1}^{\infty} \widehat{Q}_j$ converge in the strong operator topology and, moreover,

$$\sum_{n=1}^{\infty} \|P_{n,0} - \sum_{\lambda_j \in \Delta_n} \widehat{P}_j\|^2 < \infty, \quad \sum_{n=1}^{\infty} \|Q_{n,0} - \sum_{\lambda_j \in \Delta_n} \widehat{Q}_j\|^2 < \infty. \quad (2.34)$$

Proof. Convergence of the series $\sum_{j=1}^{\infty} \widehat{P}_j$ and $\sum_{j=1}^{\infty} \widehat{Q}_j$ follows from Lemma 2.11. We prove the first inequality in (2.34); the proof of the second one is similar. Recall that $P_{n,0} = \Psi_0(\pi n) \Psi_0^*(\pi n)$ and that (see (2.17))

$$\sum_{\lambda_j \in \Delta_n} \Psi_0(\pi n) \alpha_j \Psi_0^*(\pi n) = P_{n,0} - \Psi_0(\pi n) \beta_n \Psi_0^*(\pi n).$$

Using this and the relation

$$\widehat{P}_j - \Psi_0(\pi n) \alpha_j \Psi_0^*(\pi n) = \Psi_{\tau}(\lambda_j) \alpha_j [\Psi_{\tau}^*(\lambda_j) - \Psi_0^*(\pi n)] + [\Psi_{\tau}(\lambda_j) - \Psi_0(\pi n)] \alpha_j \Psi_0^*(\pi n),$$

one concludes that

$$\|P_{n,0} - \sum_{\lambda_j \in \Delta_n} \widehat{P}_j\|^2 \leq C_1 \|\beta_n\|^2 + C_2 \sum_{\lambda_j \in \Delta_n} \|\Psi_{\tau}(\lambda_j) - \Psi_0(\pi n)\|^2,$$

where C_1 and C_2 are positive constants independent of $n \in \mathbb{N}$. The result now follows from (2.30) and assumption (A_1) . $\square$

Let $\tau \in \mathbf{L}_2$ and let $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ satisfy condition (A_1) . We denote by $\mathcal{U}_{\mathbf{a},\tau}^e$ and $\mathcal{U}_{\mathbf{a},\tau}^o$ the operators given by the formulas

$$\mathcal{U}_{\mathbf{a},\tau}^e := \sum_{j=0}^{\infty} \Psi_{\tau}(\lambda_j) \alpha_j \Psi_{\tau}^*(\lambda_j), \quad \mathcal{U}_{\mathbf{a},\tau}^o := \sum_{j=1}^{\infty} \Phi_{\tau}(\lambda_j) \alpha_j \Phi_{\tau}^*(\lambda_j). \quad (2.35)$$

Remark 2.13 *For every $\tau \in \text{Re } \mathbf{L}_2$ the operators $\mathcal{U}_{\mathbf{a},\tau}^e$ and $\mathcal{U}_{\mathbf{a},\tau}^o$ are nonnegative. In addition, in view of Theorem 2.4 for every $\tau \in \text{Re } \mathbf{L}_2$ we get the equality*

$$\mathcal{U}_{\mathbf{a},\tau}^e = \mathcal{I} = \mathcal{U}_{\mathbf{a},\tau}^o. \quad (2.36)$$

Lemma 2.14 *Assume that $\tau \in \mathbf{L}_2$ and that $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ satisfies condition (A_1) . Then the following equivalences hold:*

$$(\mathcal{U}_{\mathbf{a},\tau}^e > 0) \iff (A_3), \quad (\mathcal{U}_{\mathbf{a},\tau}^o > 0) \iff (A_4). \quad (2.37)$$

Proof. Taking into account the relations (2.8), we obtain that

$$\mathcal{U}_{\mathbf{a},\tau}^e = (\mathcal{I} + \mathcal{K}_{\tau,N})\mathcal{U}_{\mathbf{a},0}^e(\mathcal{I} + \mathcal{K}_{\tau,N}^*), \quad \mathcal{U}_{\mathbf{a},\tau}^o = (\mathcal{I} + \mathcal{K}_{\tau,D})\mathcal{U}_{\mathbf{a},0}^o(\mathcal{I} + \mathcal{K}_{\tau,D}^*). \quad (2.38)$$

Since the operators $\mathcal{I} + \mathcal{K}_{\tau,N}$ and $\mathcal{I} + \mathcal{K}_{\tau,D}$ are homeomorphisms of the space L_2 , it is enough to prove equivalences (2.37) only for $\tau = 0$. Set

$$\mathcal{X} = \{d \cos \lambda_j x \mid j \in \mathbb{Z}_+, d \in \text{Ran } \alpha_j\}, \quad \mathcal{Y} = \{d \sin \lambda_j x \mid j \in \mathbb{N}, d \in \text{Ran } \alpha_j\}$$

and observe that conditions (A_3) and (A_4) are equivalent to completeness of the sets $\mathcal{X}$ and $\mathcal{Y}$ respectively. Since, in view of (2.7),

$$\Psi_0^*(\lambda)f = \sqrt{2} \int_0^1 (\cos \lambda x) f(x) dx, \quad \Phi_0^*(\lambda)f = \sqrt{2} \int_0^1 (\sin \lambda x) f(x) dx,$$

we conclude that

$$\ker \mathcal{U}_{\mathbf{a},0}^e = \bigcap_{j \in \mathbb{Z}_+} \ker \alpha_j \Psi_0^*(\lambda_j) = \mathcal{X}^\perp, \quad \ker \mathcal{U}_{\mathbf{a},0}^o = \bigcap_{j \in \mathbb{N}} \ker \alpha_j \Phi_0^*(\lambda_j) = \mathcal{Y}^\perp.$$

This justifies the equivalences of (2.37). $\square$

Proof of Theorem 2.9 Let $\tau \in \text{Re } \mathbf{L}_2$ and $\mathbf{a}_\tau = ((\lambda_j(\tau), \alpha_j(\tau)))_{j \in \mathbb{Z}_+}$. Then (A_1) holds by Theorem 2.5, while (A_3) and (A_4) are satisfied in view of (2.36) and Lemma 2.14. Next, in virtue of Lemma 2.12,

$$\sum_{n=0}^{\infty} \|P_{n,0} - \sum_{\lambda_j \in \Delta_n} P_{j,\tau}\|^2 < \infty.$$

Applying now Lemma B.1, we conclude that there exists $N_0 \in \mathbb{N}$ such that

$$\sum_{n=0}^N \sum_{\lambda_j \in \Delta_n} \text{rank } P_{j,\tau} = \sum_{n=0}^N \text{rank } P_{n,0}, \quad N \geq N_0. \quad (2.39)$$

It follows from (2.9) and (2.16) that $\text{rank } P_{j,\tau} = \text{rank } \alpha_j$ and $\text{rank } P_{n,0} = r$ for all $j, n \in \mathbb{Z}_+$. Also, $\text{rank } \alpha_0 = r$ as $\alpha_0 > 0$. Together with (2.39) this justifies (A_2) . The proof is complete. $\square$

3. The Krein accelerant

3.1. Proof of Theorem 1.7

Let H be an even function belonging to $L_2((-1, 1), M_r)$. We denote by $\mathcal{H}$ an operator in L_2 given by

$$(\mathcal{H}f)(x) := \int_0^1 H(x-t)f(t) dt. \quad (3.1)$$

Set $\mathcal{H}^a := \chi_a \mathcal{H} \chi_a$, $a \in [0, 1]$, where χ_a is an operator in L_2 of multiplication by the indicator of the interval $(0, a]$, i.e.,

$$(\chi_a f)(x) = \begin{cases} f(x), & \text{if } x \in (0, a], \\ 0, & \text{if } x \in (a, 1). \end{cases}$$

Remark 3.1 Taking into account Definition 1.5, it is easy to see that the following equivalences take place:

$$\begin{aligned} H \in \mathfrak{H}_2 &\iff \forall a \in [0, 1] \quad \ker(\mathcal{I} + \mathcal{H}^a) = \{0\}; \\ H \in \mathfrak{H}_2 &\iff H^* \in \mathfrak{H}_2; \\ H \in \text{Re } \mathfrak{H}_2 &\iff (\mathcal{I} + \mathcal{H}) > 0. \end{aligned} \tag{3.2}$$

Definition 1.5 also implies that the set $\mathfrak{H}_2$ is open and, therefore, the set $\mathfrak{H}_2 \cap C^1([-1, 1], M_r)$ is dense everywhere in $\mathfrak{H}_2$.

Suppose that $H \in \mathfrak{H}_2$. Then for every $a \in [0, 1]$ the operator $\mathcal{I} + \mathcal{H}^a$ is invertible in the algebra $\mathcal{B}(L_2)$ of bounded linear operators acting in L_2 . Since the operator $\mathcal{H}^a$ depends continuously on $a \in [0, 1]$, the mapping $[0, 1] \ni a \mapsto (\mathcal{I} + \mathcal{H}^a)^{-1} \in \mathcal{B}(L_2)$ is continuous. Denote by $\Gamma_{a,H}$ the kernel of the integral operator $-(\mathcal{I} + \mathcal{H}^a)^{-1} \mathcal{H}^a$. Since $\mathcal{H}$ belongs to the class of the Hilbert–Schmidt operators, the mapping

$$[0, 1] \times \mathfrak{H}_2 \ni (a, H) \mapsto \Gamma_{a,H} \in L_2((0, 1)^2, M_r) \tag{3.3}$$

is continuous.

It can be easily seen that, for all $(x, t) \in (0, 1)^2$ and $H \in \mathfrak{H}_2$,

$$[\Gamma_{a,H}(x, t)]^* = \Gamma_{a,H^*}(t, x) = \Gamma_{a,H^*}(x, t). \tag{3.4}$$

For every $H \in \mathfrak{H}_2$ and every $(x, t) \in \Omega$ we put

$$\tilde{R}_H(x, t) = \int_0^x H(x-y)H(y-t) dy + \int_0^x \int_0^x H(x-u)\Gamma_{x,H}(u, v)H(v-t) dv du. \tag{3.5}$$

Since the functions (3.3) are continuous and the shift $t \mapsto H(\cdot - t)$ acts continuously in L_2 -topology, one concludes that $\tilde{R}_H \in C(\Omega, M_r)$ and that the mapping

$$\mathfrak{H}_2 \ni H \mapsto \tilde{R}_H \in C(\Omega, M_r) \tag{3.6}$$

is continuous.

Lemma 3.2 Let $H \in \mathfrak{H}_2$. Then the function

$$R_H(x, t) := \begin{cases} \tilde{R}_H(x, t) - H(x-t), & 0 \leq t \leq x \leq 1, \\ 0, & 0 \leq x < t \leq 1, \end{cases} \tag{3.7}$$

belongs to G_2^+ and equation (1.11) has a unique solution in the class $L_2(\Omega, M_r)$. Moreover, the mapping $\mathfrak{H}_2 \ni H \mapsto R_H \in G_2^+$ is continuous.

Proof. That R_H belongs to G_2^+ and the mapping $H \mapsto R_H$ is continuous easily follows from continuity of the mapping (3.6). Taking into account the definition of $\Gamma_{a,H}$, we obtain that

$$H(x-t) + \Gamma_{x,H}(x, t) + \int_0^x \Gamma_{x,H}(x, \xi)H(\xi-t) d\xi = 0, \quad (x, t) \in \Omega.$$

Straightforward calculations give

$$\Gamma_{x,H}(x, t) = \tilde{R}_H(x, t) - H(x-t) = R_H(x, t), \quad (x, t) \in \Omega. \tag{3.8}$$

Thus R_H is a solution of equation (1.11). Uniqueness follows from Lemma A.3. $\square$

The proposition below follows from the results of [12] (see also [36]).

Proposition 3.3 *Suppose that $H \in \mathfrak{H}_2 \cap C([-1, 1], M_r)$. Then, for every $x, t \in [0, a]$,*

$$\frac{\partial \Gamma_{a,H}}{\partial a}(x, t) = \Gamma_{a,H}(x, a) \Gamma_{a,H}(a, t), \quad \Gamma_{a,H}(x, t) = \Gamma_{a,H}(a - x, a - t). \quad (3.9)$$

Now we study the properties of the mapping Θ defined by (1.12).

Lemma 3.4 *(i) The mapping $\Theta : \mathfrak{H}_2 \rightarrow \mathbf{L}_2$ is continuous and $\Theta(H^*) = [\Theta(H)]^*$ for every $H \in \mathfrak{H}_2$;*

(ii) if $H \in \mathfrak{H}_2 \cap C^1([-1, 1], M_r)$, then $R_H \in C^1(\Omega, M_r)$ and

$$\frac{\partial}{\partial x}[R_H(x, x - t)] - R_H(x, 0)R_H(x, t) = 0, \quad (x, t) \in \Omega. \quad (3.10)$$

Proof. *(i)* Continuity of the mapping Θ easily follows from definition and the continuity of the mapping $H \mapsto R_H$ (see Lemma 3.2). By virtue of (3.8) and the relation

$$\Gamma_{x,H}(x, \xi) = -H(x - \xi) - \int_0^x H(x - u) \Gamma_{x,H}(u, \xi) du, \quad (x, \xi) \in \Omega,$$

we obtain that

$$\Theta(H)(x) = H(x) - \int_0^x H(x - \xi) H(\xi) d\xi - \int_0^x \int_0^x H(x - u) \Gamma_{x,H}(u, v) H(v) du dv.$$

Now assume that, in addition, $H \in C([-1, 1], M_r)$; using then (3.4) and the second equality in (3.9), we get that

$$\begin{aligned} [\Theta(H)]^*(x) &= H^*(x) - \int_0^x H^*(\xi) H^*(x - \xi) d\xi \\ &\quad - \int_0^x \int_0^x H^*(v) \Gamma_{x,H^*}(v, u) H^*(x - u) du dv \\ &= H^*(x) - \int_0^x H^*(x - y) H^*(y) dy \\ &\quad - \int_0^x \int_0^x H^*(x - y) \Gamma_{x,H^*}(y, t) H^*(t) dy dt \\ &= \Theta(H^*)(x) \end{aligned}$$

as claimed. Since the set $\mathfrak{H}_2 \cap C([-1, 1], M_r)$ is dense in $\mathfrak{H}_2$ and Θ is continuous, the equality $\Theta(H^*) = [\Theta(H)]^*$ holds for all $H \in \mathfrak{H}_2$.

(ii) Let $H \in \mathfrak{H}_2 \cap C^1([-1, 1], M_r)$ and $R = R_H$, so that

$$R(x, t) + H(x - t) + \int_0^x R(x, \xi) H(\xi - t) d\xi = 0, \quad (x, t) \in \Omega. \quad (3.11)$$

It follows from Proposition 3.3 that the function $a \mapsto \Gamma_{a,H}(u, v)$ is continuously differentiable for $a \geq \max\{u, v\}$. Therefore, taking into account (3.9), (3.5), and (3.7), we obtain $R \in C^1(\Omega, M_r)$. Since R is a solution of the equation (1.11) and the function H is even, one derives the equality

$$R(x, x - t) + H(t) + \int_0^x R(x, x - \xi) H(\xi - t) d\xi = 0, \quad (x, t) \in \Omega. \quad (3.12)$$

Now differentiate (3.12) in the variable x to get

$$\frac{\partial}{\partial x}[R(x, x-t)] + R(x, 0)H(x-t) + \int_0^x \frac{\partial}{\partial x}[R(x, x-\xi)]H(\xi-t) d\xi = 0. \quad (3.13)$$

Multiplying both sides of equality (3.11) by $R(x, 0)$ from the left, we obtain

$$R(x, 0)R(x, t) + R(x, 0)H(x-t) + \int_0^x R(x, 0)R(x, \xi)H(\xi-t) d\xi = 0.$$

Subtracting (3.13) from the above equality, we find that for the function

$$F(x, t) := \frac{\partial}{\partial x}[R(x, x-t)] - R(x, 0)R(x, t)$$

the following relation holds for all $(x, t) \in \Omega$:

$$F(x, t) + \int_0^x F(x, \xi)H(\xi-t) d\xi = 0.$$

Lemma A.3 now yields $F \equiv 0$, and the proof is complete. $\square$

Proof of Theorem 1.7 Assume that $H \in \mathfrak{H}_2$ and set

$$Q_{o,H}(x, t) := \frac{1}{2} \left[R_H\left(x, \frac{x+t}{2}\right) - R_H\left(x, \frac{x-t}{2}\right) \right],$$

$$Q_{e,H}(x, t) := \frac{1}{2} \left[R_H\left(x, \frac{x+t}{2}\right) + R_H\left(x, \frac{x-t}{2}\right) \right],$$

where $(x, t) \in \Omega$. We extend the functions $Q_{o,H}$ and $Q_{e,H}$ onto the square $(0, 1)^2$ by setting $Q_{o,H} \equiv 0$ and $Q_{e,H} \equiv 0$ in $\Omega^- = [0, 1]^2 \setminus \Omega$. It follows from (3.7) and the continuity of the mapping (3.6) that $Q_{o,H}$ and $Q_{e,H}$ belong to G_2 and that the mappings

$$\mathfrak{H}_2 \ni H \mapsto Q_{o,H} \in G_2, \quad \mathfrak{H}_2 \ni H \mapsto Q_{e,H} \in G_2 \quad (3.14)$$

are continuous. For $x \in [0, 1]$ and $\lambda \in \mathbb{C}$, we consider the functions

$$\varphi(x, \lambda) := \sin \lambda x I + \int_0^x (\sin \lambda t) Q_{o,H}(x, t) dt,$$

$$\psi(x, \lambda) := \cos \lambda x I + \int_0^x (\cos \lambda t) Q_{e,H}(x, t) dt.$$

Straightforward transformations give

$$\varphi(x, \lambda) = \sin \lambda x I + \int_0^x (\sin \lambda(x-2s)) R(x, x-s) ds,$$

$$\psi(x, \lambda) = \cos \lambda x I + \int_0^x (\cos \lambda(x-2s)) R(x, x-s) ds. \quad (3.15)$$

Suppose that $H \in \mathfrak{H}_2 \cap C^1([-1, 1], M_r)$. According to Lemma 3.4, the function R_H belongs to $C^1(\Omega, M_r)$ and equality (3.10) holds. Using (3.15) and (3.10), we arrive at the relations

$$\left(\frac{d}{dx} - \tau \right) \varphi(x, \lambda) = \lambda \psi(x, \lambda), \quad \left(\frac{d}{dx} + \tau \right) \psi(x, \lambda) = -\lambda \varphi(x, \lambda),$$

where $\tau = \Theta(H)$. Since $\varphi(0, \lambda) = 0$ and $\psi(0, \lambda) = I$, the functions φ and ψ are solutions of problems (1.3) and (1.4). In view of Theorem 2.1 this establishes the relations

$$Q_{o,H} = K_{\tau,D}, \quad Q_{e,H} = K_{\tau,N}, \quad \tau = \Theta(H) \quad (3.16)$$

for all $H \in \mathfrak{H}_2 \cap C^1([-1, 1], M_r)$. Since the set $\mathfrak{H}_2 \cap C^1([-1, 1], M_r)$ is dense everywhere in $\mathfrak{H}_2$ and the mappings Θ , (3.14), $\mathbf{L}_2 \ni \tau \rightarrow K_{\tau,D} \in G_2^+$, and $\mathbf{L}_2 \ni \tau \rightarrow K_{\tau,N} \in G_2^+$ are continuous (see Theorem 2.1), it follows that equalities (3.16) hold for all $H \in \mathfrak{H}_2$. The proof is complete. $\square$

3.2. Proof of Theorem 1.8.

Consider the integral equation

$$P(x, x-s) + \tau(s) + \int_s^x \tau(u)P(u, s) du = 0, \quad (x, s) \in \Omega, \quad (3.17)$$

where τ is a known function from $\mathbf{L}_2$, and P is an unknown function from the class G_2^+ .

Lemma 3.5 *The equation (3.17) has at most one solution. If $H \in \mathfrak{H}_2 \cap C^1([-1, 1], M_r)$ and $\tau = \Theta(H)$, then the solution R_H of the Krein equation (1.11) is also a solution of the equation (3.17).*

Proof. To prove uniqueness, it is enough to show that the corresponding homogeneous equation

$$F(x, x-s) = - \int_s^x \tau(u)F(u, s) du, \quad (x, s) \in \Omega, \quad (3.18)$$

has only zero solution in G_2^+ . Let a function $F \in G_2^+$ be a solution of (3.18) and set

$$g(x) := \int_0^x \|F(x, s)\| ds, \quad x \in [0, 1].$$

Then g is continuous on $[0, 1]$ and, in view of (3.18),

$$g(x) = \int_0^x \|F(x, x-s)\| ds \leq \int_0^x \|\tau(u)\| \int_0^u \|F(u, s)\| ds du \leq \int_0^x \|\tau(u)\| g(u) du.$$

Now Gronwall's inequality implies that $g \equiv 0$, whence $F = 0$.

Let $H \in \mathfrak{H}_2$ and $\tau = \Theta(H)$. Since $\tau = \Theta(H) = -R_H(\cdot, 0)$, we get by virtue of equality (3.10) that

$$R_H(x, x-t) + \tau(t) + \int_t^x \tau(\xi)R_H(\xi, t) d\xi = 0, \quad (x, t) \in \Omega.$$

Thus R_H is a solution of equation (3.17). $\square$

Lemma 3.6 *For every $\tau \in \mathbf{L}_2$ equation (3.17) has a unique solution P_τ in G_2^+ and the mapping $\mathbf{L}_2 \ni \tau \mapsto P_\tau \in G_2^+$ is continuous. Moreover, if $\tau \in C([0, 1], M_r)$, then P_τ is continuous in Ω , the function $(x, t) \mapsto P_\tau(x, x-t)$ has continuous partial derivative in the variable x , and*

$$\frac{\partial}{\partial x}[P_\tau(x, x-t)] = -\tau(x)P_\tau(x, t), \quad (x, t) \in \Omega. \quad (3.19)$$

Proof. For every $n \in \mathbb{N}$ and $(x, s) \in \Omega$, we set

$$P_{\tau,1}(x, x-s) := \tau(s), \quad P_{\tau,n+1}(x, x-s) := \int_s^x \tau(u) P_{\tau,n}(u, s) du. \quad (3.20)$$

It follows by induction that the recurrence relation (3.20) yields the equality

$$P_{\tau,n+1}(x, s) = \int_{\Pi_n^*(x,s)} \tau(x-s+\xi_n(y)) \tau(y_1) \cdots \tau(y_n) dy_1 \cdots dy_n, \quad (3.21)$$

where, for $n \in \mathbb{N}$ and $(x, s) \in \Omega$, we set $\xi_n(y) := \sum_{j=1}^n (-1)^{j+1} y_j$ and

$$\Pi_n^*(x, s) := \{y = (y_1, \dots, y_n) \in \mathbb{R}^n \mid 0 \leq y_n \leq y_{n-1} \leq \dots \leq y_1 \leq x-s+\xi_n(y) \leq x\}.$$

We extend the functions $P_{\tau,n}$, $n \in \mathbb{N}$, onto the square $[0, 1] \times [0, 1]$ by setting $P_{\tau,n} \equiv 0$ in Ω^- . Since the function τ belongs to $\mathbf{L}_2$, for any $n \in \mathbb{N}$ the functions $P_{\tau,n+1}$ are continuous in the square $[0, 1]^2$. Taking into account (3.20) and using the Cauchy–Bunyakovski inequality and Fubini’s theorem, we find that for every $n \in \mathbb{N}$ and $x \in [0, 1]$ it holds

$$\begin{aligned} \|P_{\tau,n+1}(x, \cdot)\|_{\mathbf{L}_2}^2 &= \int_0^1 \|P_{\tau,n+1}(x, s)\|^2 ds \\ &\leq \frac{1}{n!} \int_0^1 \int_{\Pi_n^*(x,s)} \|\tau(x-s+\xi_{n-1}(y))\|^2 \|\tau(y_1)\|^2 \cdots \|\tau(y_n)\|^2 dy_1 \cdots dy_n ds \\ &\leq \frac{1}{n!} \int_{\Pi_{n+1}} \|\tau(t_1)\|^2 \cdots \|\tau(t_{n+1})\|^2 dt_1 \cdots dt_{n+1} \\ &= \frac{1}{(n!)(n+1)!} \left(\int_0^1 \|\tau(\xi)\|^2 d\xi \right)^{n+1}, \end{aligned}$$

where $\Pi_n = \{t := (t_1, \dots, t_n) \in \mathbb{R}^n \mid 0 \leq t_n \leq \dots \leq t_1 \leq 1\}$. Therefore

$$\|P_{\tau,n+1}(x, \cdot)\|_{\mathbf{L}_2} \leq (n!)^{-1} \|\tau\|_{\mathbf{L}_2}^{n+1}.$$

Similarly we obtain that

$$\|P_{\tau,n+1}(\cdot, x)\|_{\mathbf{L}_2} \leq (n!)^{-1} \|\tau\|_{\mathbf{L}_2}^{n+1}.$$

Now, taking into account continuity of the functions $P_{\tau,n+1}$, we conclude that $P_{\tau,n+1}$ belongs to G_2^+ , and, moreover,

$$\|P_{\tau,n+1}\|_{G_2} \leq (n!)^{-1} \|\tau\|_{\mathbf{L}_2}^{n+1}, \quad n \in \mathbb{N}. \quad (3.22)$$

Taking into account (3.20) and (3.22), we see that the function

$$P_\tau := \sum_{n=1}^{\infty} (-1)^n P_{\tau,n} \quad (3.23)$$

is a solution of equation (3.17). Uniqueness of solution of equation (3.17) follows from Lemma 3.5.

Next we prove continuous dependence of P_τ on τ . It follows from (3.22) and (3.23) that it is sufficient to prove continuity of the mappings

$$\mathbf{L}_2 \ni \tau \mapsto P_{\tau,n} \in G_2^+, \quad n \in \mathbb{N}. \quad (3.24)$$

Let $\tau_1, \tau_2 \in \mathbf{L}_2$. Using (3.21) and the inequality

$$\left| \prod_{k=1}^n a_k - \prod_{k=1}^n b_k \right|^2 \leq n \sum_{k=1}^n |a_k - b_k|^2 \prod_{j \neq k} (|a_j| + |b_j|)^2,$$

valid for all $(a_k)_{k=1}^n$ and $(b_k)_{k=1}^n$ in $\mathbb{C}^n$, and reasoning similarly as when deriving the estimate (3.22), we obtain the estimate

$$\|P_{\tau_1, n+1} - P_{\tau_2, n+1}\|_{G_2^+}^2 \leq \left(\frac{n+1}{n!} \right)^2 \|\tau_1 - \tau_2\|_{\mathbf{L}_2}^2 \left(\|\tau_1\|_{\mathbf{L}_2} + \|\tau_2\|_{\mathbf{L}_2} \right)^{2n},$$

which yields continuity of the mappings (3.24).

Finally, assume that $\tau \in C([0, 1], M_r)$. Since the function P_τ verifies (3.17), P_τ is continuous, and the function $(x, s) \mapsto P_\tau(x, x - s)$ is absolutely continuous in x . Differentiation of (3.17) in the variable x leads to (3.19). The proof is complete. $\square$

Proof of Theorem 1.8 For any $\tau \in \mathbf{L}_2$ we denote by $\mathcal{P}_\tau$ the integral operator with kernel P_τ . It follows from Lemma 3.6 that $\mathcal{P}_\tau$ belongs to $\mathfrak{G}_2^+$ and that the mapping $\mathbf{L}_2 \ni \tau \mapsto \mathcal{P}_\tau \in \mathfrak{G}_2^+$ is continuous. Hence in view of Lemma A.4 we obtain that the mapping Υ that acts from $\mathbf{L}_2$ to $L_2((-1, 1), M_r)$ via

$$[\Upsilon(\tau)](x) = [(\mathcal{I} + \mathcal{P}_\tau)^{-1}\tau](|x|), \quad x \in (-1, 1),$$

is also continuous. Next we show that the mapping Υ is the left inverse of Θ , i.e., that the relation

$$\Upsilon(\Theta(H)) = H \tag{3.25}$$

holds for all $H \in \mathfrak{H}_2$. Since the set $\mathfrak{H}_2 \cap C^1([-1, 1], M_r)$ is dense everywhere in $\mathfrak{H}_2$ and the mappings Θ and Υ are continuous, it is enough to prove equality (3.25) for $H \in \mathfrak{H}_2 \cap C^1([-1, 1], M_r)$. Assume H is such and set $\tau = \Theta(H)$. Then it follows from Lemmas 3.5 and 3.6 that $P_\tau = R_H$. Therefore, recalling (1.12), we obtain that

$$\tau(x) = [\Theta(H)](x) = H(x) + \int_0^x P_\tau(x, \xi) H(\xi) d\xi, \quad x \in (0, 1),$$

and, therefore, $\Upsilon(\Theta(H)) = \Upsilon(\tau) = H$.

It remains to show that Υ is also the right inverse of Θ , i.e., that, for every $\tau \in \mathbf{L}_2$,

$$\Upsilon(\tau) \in \mathfrak{H}_2 \quad \text{and} \quad \Theta(\Upsilon(\tau)) = \tau. \tag{3.26}$$

Continuity of the mappings Θ and Υ implies that it is enough to prove (3.26) for continuous τ . In this case for $H := \Upsilon(\tau)$ we get the relation

$$\tau(x) = H(x) + \int_0^x P_\tau(x, \xi) H(\xi) d\xi, \quad x \in [0, 1]. \tag{3.27}$$

According to Lemma 3.6, the function P_τ is continuous in Ω , and, therefore, the function H is continuous as well. Consider a function F defined for $(x, t) \in \Omega$ by

$$F(x, t) := P_\tau(x, x - t) + H(t) + \int_0^x P_\tau(x, x - \xi) H(\xi - t) d\xi.$$

Then the function F is continuous in Ω , has a continuous partial derivative in the variable x by virtue of Lemma 3.6, and

$$\frac{\partial}{\partial x}[F(x, x-t)] = -\tau(x)F(x, t), \quad (x, t) \in \Omega.$$

Since

$$F(t, 0) = -\tau(t) + H(t) + \int_0^t P_\tau(t, \xi)H(\xi) d\xi = 0, \quad t \in [0, 1],$$

we obtain that

$$F(x, x-t) = -\int_t^x \tau(\xi)F(\xi, t) d\xi, \quad t \leq x \leq 1.$$

However, as it was shown in the proof of Lemma 3.5, the last equality is only possible for $F \equiv 0$. Therefore P_τ is a solution of the Krein equation (1.11) with $H = \Upsilon(\tau)$. Now Proposition A.2 implies that H is an accelerant; moreover, $\Theta(\Upsilon(\tau)) = \tau$ in view of (3.27). Finally, the equality $\Theta(H^*) = [\Theta(H)]^*$ follows from Lemma 3.4 (i). The proof is complete. $\square$

3.3. The operators $\mathcal{H}_o$ and $\mathcal{H}_e$

For any $H \in L_2((-1, 1), M_r)$ we denote by $\mathcal{H}_o$ and $\mathcal{H}_e$ integral operators that act in L_2 by the formulas

$$(\mathcal{H}_o f)(x) := \int_0^1 H_o(x, t)f(t) dt, \quad (\mathcal{H}_e f)(x) := \int_0^1 H_e(x, t)f(t) dt, \quad (3.28)$$

where

$$\begin{aligned} H_o(x, t) &:= \frac{1}{2} \left[H\left(\frac{x-t}{2}\right) - H\left(\frac{x+t}{2}\right) \right], \\ H_e(x, t) &:= \frac{1}{2} \left[H\left(\frac{x-t}{2}\right) + H\left(\frac{x+t}{2}\right) \right]. \end{aligned}$$

Proposition 3.7 *Let $H \in \mathfrak{H}_2$ and $\tau = \Theta(H)$. Then*

$$(\mathcal{I} + \mathcal{K}_{\tau, N})(\mathcal{I} + \mathcal{H}_e)(\mathcal{I} + \mathcal{K}_{\tau^*, N}^*) = \mathcal{I} = (\mathcal{I} + \mathcal{K}_{\tau, D})(\mathcal{I} + \mathcal{H}_o)(\mathcal{I} + \mathcal{K}_{\tau^*, D}^*). \quad (3.29)$$

Proof. We shall prove the first equality in (3.29); the second one is proved similarly. It follows from the Krein equation (1.11) that

$$\begin{aligned} R_H\left(x, \frac{x-t}{2}\right) + H\left(\frac{x+t}{2}\right) + \int_0^x R_H(x, \xi)H\left(\xi - \frac{x-t}{2}\right) d\xi &= 0, \\ R_H\left(x, \frac{x+t}{2}\right) + H\left(\frac{x-t}{2}\right) + \int_0^x R_H(x, \xi)H\left(\xi - \frac{x+t}{2}\right) d\xi &= 0 \end{aligned}$$

for $(x, t) \in \Omega$. Combining these relations and taking into account Theorem 1.7 proved above, we arrive at the equality

$$K_{\tau, N}(x, t) + H_e(x, t) + \int_0^x K_{\tau, N}(x, \xi)H_e(\xi, t) d\xi = 0 \quad (3.30)$$

for all $(x, t) \in \Omega$. By virtue of Lemma 3.4, $\Theta(H^*) = [\Theta(H)]^* = \tau^*$. Thus, according to (3.30), we find that for $(x, t) \in \Omega$ it holds

$$K_{\tau^*, N}(x, t) + H_e^*(x, t) + \int_0^x K_{\tau^*, N}(x, \xi) H_e^*(\xi, t) d\xi = 0. \quad (3.31)$$

Equalities (3.30), (3.31), and Theorem A.1 now lead to the relation

$$\mathcal{I} + \mathcal{H}_e = (\mathcal{I} + \mathcal{K}_{\tau, N})^{-1}(\mathcal{I} + \mathcal{K}_{\tau^*, N}^*)^{-1},$$

which yields the first equality in (3.29). The lemma is proved. $\square$

Proposition 3.8 *If $\ker(\mathcal{I} + \mathcal{H}_e) = \{0\}$, then $\mathcal{I} + \mathcal{H}_e$ is a bijection of W_2^1 .*

Proof. It is enough to prove that $\mathcal{H}_e$ maps W_2^1 to itself and that the operator $\mathcal{H}_e : W_2^1 \rightarrow W_2^1$ is compact. If $H \in C^1([-1, 1], M_r)$, then integration by parts gives

$$(\mathcal{H}_e f)' = \mathcal{H}_e f' - H_o(\cdot, 1)f(1), \quad f \in W_2^1. \quad (3.32)$$

Using the fact that the operator of differentiation considered on the domain W_2^1 is closed in L_2 and that the space $C^1([-1, 1], M_r)$ is dense in $L_2((-1, 1), M_r)$, we conclude that the equality (3.32) holds also for an arbitrary function $H \in L_2((-1, 1), M_r)$. The equality (3.32) implies that the operator $\mathcal{H}_e$ maps the space W_2^1 to itself; moreover,

$$\|\mathcal{H}_e\|_{W_2^1 \rightarrow W_2^1} \leq C \|H\|_{L_2((-1, 1), M_r)}, \quad (3.33)$$

where C is a constant independent of H . If the function H is a trigonometric polynomial, then $\mathcal{H}_e$ is an operator of finite rank. Approximating an arbitrary function $H \in L_2((-1, 1), M_r)$ by a sequence of trigonometric polynomials and taking into account (3.33), we conclude that the operator $\mathcal{H}_e : W_2^1 \rightarrow W_2^1$ is compact. $\square$

4. The inverse spectral problem

4.1. Proof of Theorem 1.6

We shall base the proof of Theorem 1.6 on the following two lemmas.

Lemma 4.1 *Let for a sequence $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ condition (A_1) hold and $\nu = \nu^{\mathbf{a}}$. Then the limit (1.10) exists in the topology of the space $L_2((-1, 1), M_r)$. Moreover, the function $H = H_\nu$ is even and hermitian, and for the operators $\mathcal{H}_e$ and $\mathcal{H}_o$ the following relations hold:*

$$\mathcal{I} + \mathcal{H}_e = \mathcal{U}_{\mathbf{a}, 0}^e, \quad \mathcal{I} + \mathcal{H}_o = \mathcal{U}_{\mathbf{a}, 0}^o. \quad (4.1)$$

Proof. In view of (1.7) and (1.8) we have to show that the series

$$\sum_{n=1}^{\infty} \left[\cos(2\pi n x) I - \sum_{\lambda_j \in \Delta_n} \cos(2\lambda_j x) \alpha_j \right] \quad (4.2)$$

converges in $L_2((-1, 1), M_r)$. Let $\tilde{\lambda}_j$ and β_n be given by (2.17). Then

$$\cos 2\lambda_j x = \cos 2\pi n x - 2x \tilde{\lambda}_j \sin 2\pi n x - \vartheta_{n,j}(x), \quad n \in \mathbb{N}, \quad \lambda_j \in \Delta_n,$$

where

$$\vartheta_{n,j}(x) := (1 - \cos 2x\tilde{\lambda}_j) \cos 2\pi nx - (2\tilde{\lambda}_j x - \sin 2\tilde{\lambda}_j x) \sin 2\pi nx.$$

This yields the equality

$$\cos(2\pi nx)I - \sum_{\lambda_j \in \Delta_n} \cos(2\lambda_j x)\alpha_j = \cos(2\pi nx)\beta_n + 2x \sin(2\pi nx)\gamma_n + \sum_{\lambda_j \in \Delta_n} \vartheta_{n,j}(x)\alpha_j,$$

in which $\gamma_n := \sum_{\lambda_j \in \Delta_n} \tilde{\lambda}_j \alpha_j$. According to condition (A_1)

$$\sum_{n=1}^{\infty} \|\beta_n\|^2 < \infty, \quad \sup_{j \in \mathbb{Z}_+} \|\alpha_j\| < \infty, \quad \sum_{j=1}^{\infty} |\tilde{\lambda}_j|^2 < \infty, \quad \sup_{n \in \mathbb{N}} \sum_{\lambda_j \in \Delta_n} 1 < \infty, \quad (4.3)$$

which, in particular, implies that $\sum_{n=1}^{\infty} \|\gamma_n\|^2 < \infty$. From the above we conclude that the series $\sum_{n=1}^{\infty} \cos(2\pi nx)\beta_n$ and $\sum_{n=1}^{\infty} \sin(2\pi nx)\gamma_n$ converge in the topology of the space $L_2((-1, 1), M_r)$. It is easy to see that

$$\max_{|s| \leq 1} |\vartheta_{n,j}(s)| \leq 6|\tilde{\lambda}_j|^2, \quad n \in \mathbb{N}, \quad \lambda_j \in \Delta_n.$$

Therefore,

$$\sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} \|\vartheta_{n,j}\alpha_j\|_{L_2((-1,1),M_r)} \leq 6 \left(\sup_{j \in \mathbb{Z}_+} \|\alpha_j\| \right) \sum_{j=1}^{\infty} |\tilde{\lambda}_j|^2 < \infty.$$

Combining the above, we conclude that the series (4.2) converges. The fact that the function H_ν is even and hermitian is obvious.

Now construct operators $\mathcal{H}_e$ and $\mathcal{H}_o$ via formulas (3.28) taking there $H = H_\nu$. We shall prove the first equality in (4.1), since the second one is proved similarly. Let

$$H_{\nu,n}(x) := \int_0^{\pi(n+1/2)} 2 \cos(2\lambda x) d(\nu - \nu_0)(\lambda), \quad x \in [-1, 1], \quad n \in \mathbb{N}.$$

We denote by $\mathcal{H}_{e,n}$ the operator in L_2 that acts via the formula

$$(\mathcal{H}_{e,n}f)(x) := \frac{1}{2} \int_0^1 \left[H_{\nu,n}\left(\frac{x-t}{2}\right) + H_{\nu,n}\left(\frac{x+t}{2}\right) \right] f(t) dt, \quad x \in [-1, 1].$$

Since $H_{\nu,n} \rightarrow H_\nu$ as $n \rightarrow \infty$, the sequence $(\mathcal{H}_{e,n})_{n \in \mathbb{N}}$ converges in the operator norm to the operator $\mathcal{H}_e$. It is easy to see that

$$\mathcal{H}_{e,n} = \sum_{\lambda_j \leq \pi(n+1/2)} \Psi_0(\lambda_j)\alpha_j\Psi_0^*(\lambda_j) - \sum_{k=0}^n P_{k,0}.$$

The required equality follows now from (2.15) and (2.35). $\square$

Lemma 4.2 *Let a sequence $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ verifies condition (A_1) . Then the following implications are true: $(A_3) \implies (A_4)$ and $[(A_4) \wedge (\lambda_0 = 0, \alpha_0 > 0)] \implies (A_3)$.*

Proof. Let conditions (A_1) and (A_3) hold. Then it follows from (4.1) and (2.37) that $\ker(\mathcal{I} + \mathcal{H}_e) = \{0\}$. Hence, taking into account Proposition 3.8, we obtain that $(\mathcal{I} + \mathcal{H}_e)W_2^1 = W_2^1$. Therefore, for an arbitrary $g \in L_2$ there exists $f \in W_2^1$ such that $g = [(\mathcal{I} + \mathcal{H}_e)f]'$. Put $c_j := \alpha_j \Psi_0^*(\lambda_j)f$, $j \in \mathbb{Z}_+$. In view of (4.1) and (2.35)

$$(\mathcal{I} + \mathcal{H}_e)f = \sum_{j=0}^{\infty} \Psi_0(\lambda_j)c_j. \quad (4.4)$$

Taking into account the relation

$$\lambda_j \Psi_0^*(\lambda_j)f = \sqrt{2} \int_0^1 (\lambda_j \cos \lambda_j t) f(t) dt = (-1)^n \sqrt{2} (\sin \tilde{\lambda}_j) f(1) - \Phi_0^*(\lambda_j) f', \quad \lambda_j \in \Delta_n,$$

the inequalities in (4.3), and Lemma 2.11, we conclude that $\sum_{j=0}^{\infty} \|\lambda_j c_j\|_{\mathbb{C}^r}^2 < \infty$. In view of Lemma 2.11 the series in (4.4) is termwise differentiable and therefore $g = - \sum_{j=1}^{\infty} \lambda_j \Phi_0(\lambda_j)c_j$. Since g is arbitrary, the system $\{d \sin \lambda_j x \mid j \in \mathbb{N}, d \in \text{Ran } \alpha_j\}$ is complete in the space L_2 , and the implication $(A_3) \implies (A_4)$ follows.

Now let the conditions (A_1) , (A_4) hold and let $\lambda_0 = 0$ and $\alpha_0 > 0$. Then $\ker(\mathcal{I} + \mathcal{H}_o) = \{0\}$ by virtue of (4.1) and (2.37). Since the operator $\mathcal{H}_o : L_2 \rightarrow L_2$ is compact, $\text{Ran}(\mathcal{I} + \mathcal{H}_o) = L_2$. Let $J : L_2 \rightarrow L_2$ be an integration operator, i.e.,

$$(Jf)(x) = \int_0^x f(t) dt.$$

The range of the operator J is everywhere dense in L_2 . Therefore the range of the operator $J(\mathcal{I} + \mathcal{H}_o)$ is also everywhere dense in L_2 . In view of (4.1) for an arbitrary $f \in L_2$ we get

$$J(\mathcal{I} + \mathcal{H}_o)f = \sum_{j=1}^{\infty} J\Phi_0(\lambda_j)\alpha_j\Phi_0^*(\lambda_j)f = \sum_{j=1}^{\infty} \frac{\sqrt{2}}{\lambda_j} d_j - \sum_{j=1}^{\infty} \frac{1}{\lambda_j} \Psi_0(\lambda_j)d_j,$$

where $d_j := \alpha_j \Phi_0^*(\lambda_j)f$. Since the matrix α_0 is nonsingular, the vector $\sum_{j=1}^{\infty} \sqrt{2}d_j/\lambda_j$ is in the range of α_0 . Therefore the closed linear hull of the system

$$\{(\cos \lambda_j x) d \mid j \in \mathbb{Z}_+, d \in \text{Ran } \alpha_j\}$$

contains the range of $J(\mathcal{I} + \mathcal{H}_o)$ and thus (A_3) holds. $\square$

Proof of Theorem 1.6 In view of Lemma 4.1 only the second part of the theorem needs to be proved. In fact, it suffices to prove that conditions (A_1) and (A_3) imply that $H \in \text{Re } \mathfrak{H}_2$.

Let therefore conditions (A_1) and (A_3) hold. It follows then from (4.1), (2.37), and Lemma 4.2 that

$$\mathcal{I} + \mathcal{H}_e > 0, \quad \mathcal{I} + \mathcal{H}_o > 0. \quad (4.5)$$

Let us show that (4.5) implies positivity of the operator $\mathcal{I} + \mathcal{H}$. Let $L_{2,e}$ and $L_{2,o}$ be subspaces of L_2 consisting of functions that are respectively even and odd with respect to $\frac{1}{2}$, i.e.,

$$L_{2,e} := \{f \in L_2 \mid f(1-x) = f(x)\}, \quad L_{2,o} := \{f \in L_2 \mid f(1-x) = -f(x)\}.$$

$L_{2,e}$ and $L_{2,o}$ are invariant subspaces of the operator $\mathcal{H}$. Consider unitary operators $\mathcal{A}_o : L_2 \rightarrow L_{2,o}$ and $\mathcal{A}_e : L_2 \rightarrow L_{2,e}$ acting via the formulas

$$(\mathcal{A}_o g)(x) = \operatorname{sgn}(2x-1)g(|2x-1|), \quad (\mathcal{A}_e g)(x) = g(|2x-1|), \quad x \in (0,1).$$

Simple calculations show that for an arbitrary $g \in L_2$ we have

$$((\mathcal{I} + \mathcal{H})\mathcal{A}_o g | \mathcal{A}_o g) = ((\mathcal{I} + \mathcal{H}_o)g | g), \quad ((\mathcal{I} + \mathcal{H})\mathcal{A}_e g | \mathcal{A}_e g) = ((\mathcal{I} + \mathcal{H}_e)g | g).$$

Taking into account (4.5) and the equality $L_{2,o} \oplus L_{2,e} = L_2$, we obtain that $\mathcal{I} + \mathcal{H} > 0$. Therefore due to (3.2) the function H belongs to $Re \mathfrak{H}_2$. $\square$

4.2. Proof of sufficiency in Theorems 1.1 and 1.2

In Section 2 the necessity parts of Theorems 1.1 and 1.2 was justified, and we still have to establish the sufficiency parts. By virtue of Lemma 4.2 it suffices to prove the following theorem.

Theorem 4.3 *Let for a sequence $\mathbf{a} = ((\lambda_j, \alpha_j))_{j \in \mathbb{Z}_+}$ conditions (A_1) – (A_3) hold and let $\nu = \nu^a$ and $\tau = \Theta(H_\nu)$. Then $\tau \in Re \mathbf{L}_2$ and $\mathbf{a} = \mathbf{a}_\tau$.*

First we prove the following lemma.

Lemma 4.4 *Let the assumptions of Theorem 4.3 hold. Then the formulas*

$$\hat{P}_j = \Psi_\tau(\lambda_j) \alpha_j \Psi_\tau^*(\lambda_j), \quad \hat{Q}_k = \Phi_\tau(\lambda_k) \alpha_k \Phi_\tau^*(\lambda_k) \quad (4.6)$$

determine complete systems $(\hat{P}_j)_{j \in \mathbb{Z}_+}$ and $(\hat{Q}_j)_{j \in \mathbb{N}}$ of pairwise orthogonal projectors.

Proof. Taking into account Proposition 3.7 and equalities (4.1) and (2.38), we arrive at the equality $\mathcal{U}_{\mathbf{a},\tau}^e = \mathcal{I} = \mathcal{U}_{\mathbf{a},\tau}^o$ for $\tau = \Theta(H_\nu)$ and $\nu = \nu^a$. Hence, by virtue of (2.35) and Lemma 2.12, we conclude that

$$\sum_{n=0}^{\infty} \sum_{\lambda_j \in \Delta_n} \hat{P}_j = \mathcal{I} = \sum_{n=1}^{\infty} \sum_{\lambda_j \in \Delta_n} \hat{Q}_j, \quad (4.7)$$

and that both series converge in the strong operator topology. We show that $(\hat{P}_j)_{j \in \mathbb{Z}_+}$ is the sequence of pairwise orthogonal projectors; the proof for $(\hat{Q}_j)_{j \in \mathbb{N}}$ is analogous.

In view of Lemma 2.12,

$$\sum_{n=1}^{\infty} \|P_{n,0} - \sum_{\lambda_j \in \Delta_n} \hat{P}_j\|^2 < \infty.$$

Thus there exists a natural N_0 such that

$$\sum_{n=N_0}^{\infty} \|P_{n,0} - \sum_{\lambda_j \in \Delta_n} \hat{P}_j\|^2 < 1. \quad (4.8)$$

Moreover, N_0 can be taken so large that (A_2) holds, i.e., that for every $N \geq N_0$ it holds

$$\sum_{n=0}^N \sum_{\lambda_j \in \Delta_n} \operatorname{rank} \alpha_j = (N+1)r. \quad (4.9)$$

Let us fix $N > N_0$ and set

$$A = \sum_{n=N+1}^{\infty} \sum_{\lambda_j \in \Delta_n} \widehat{P}_j, \quad P = \sum_{n=0}^N P_{n,0}, \quad A_k = \sum_{\lambda_j \in \Delta_k} \widehat{P}_j.$$

Since $\text{rank } \widehat{P}_j = \text{rank } \alpha_j$ for all $j \in \mathbb{Z}_+$ in view of (2.9) and (4.6), we conclude by virtue of (4.9) that, for all $k > N_0$,

$$\text{rank } A_k \leq \sum_{\lambda_j \in \Delta_k} \text{rank } \widehat{P}_j = \sum_{\lambda_j \in \Delta_k} \text{rank } \alpha_j = r = \text{rank } P_{k,0}. \quad (4.10)$$

Since $(P_{n,0})_{n \in \mathbb{Z}_+}$ is a complete sequence of pairwise orthogonal projectors, by virtue of (4.8), (4.10), and Lemma B.2 we obtain that

$$\text{codim } \text{Ran } A \geq \text{rank } P = (N+1)r. \quad (4.11)$$

It follows from (4.7), (4.9), and (4.11) that

$$\begin{aligned} A + \sum_{n=0}^N \sum_{\lambda_j \in \Delta_n} \widehat{P}_j &= \mathcal{I}, \\ \sum_{n=0}^N \sum_{\lambda_j \in \Delta_n} \text{rank } \widehat{P}_j &= \sum_{n=0}^N \sum_{\lambda_j \in \Delta_n} \text{rank } \alpha_j = (N+1)r \leq \text{codim } \text{Ran } A. \end{aligned}$$

The above relations imply by virtue of Lemma B.3 that the set $\{\widehat{P}_j : \lambda_j \leq \pi N + \pi/2\}$ is a set of pairwise orthogonal projectors. Since N was arbitrary, we conclude that $(\widehat{P}_j)_{j \in \mathbb{Z}_+}$ is a sequence of pairwise orthogonal projectors. The proof is complete. $\square$

Proof of Theorem 4.3. Let the assumptions of Theorem 4.3 hold and let $\widehat{P}_j$ and $\widehat{Q}_j$ be operators of Lemma 4.2. It follows from Theorems 1.6 and 1.8 that $\tau \in \text{Re } \mathbf{L}_2$. Theorem 4.3 will be proved if we show that

$$\text{Ran } \widehat{P}_j \subset \ker(S_\tau - \lambda_j^2 \mathcal{I}), \quad j \in \mathbb{Z}_+. \quad (4.12)$$

Indeed, (4.12) together with (4.7) imply that $\lambda_j(\tau) = \lambda_j$ for all $j \in \mathbb{Z}_+$. The last equality and (4.12) yield the inequality $P_{j,\tau} - \widehat{P}_j \geq 0$, $j \in \mathbb{Z}_+$. However, in view of (4.7) and (2.15) we have

$$\sum_{j=0}^{\infty} (P_{j,\tau} - \widehat{P}_j) = 0.$$

Thus $P_{j,\tau} - \widehat{P}_j = 0$, $j \in \mathbb{Z}_+$, and, therefore, (see (2.16) and (4.6))

$$\Psi_\tau(\lambda_j)[\alpha_j(\tau) - \alpha_j]\Psi_\tau^*(\lambda_j) = 0, \quad j \in \mathbb{Z}_+.$$

Now by virtue of (2.9) we obtain the equality $\alpha_j(\tau) = \alpha_j$, $j \in \mathbb{Z}_+$, i.e., the equality $\mathbf{a} = \mathbf{a}_\tau$.

It thus indeed only remains to prove (4.12). In view of (2.7) and (2.1) it is enough to show that $\varphi(1, \lambda_j, \tau)\alpha_j = 0$, $j \in \mathbb{Z}_+$. Let $j, k \in \mathbb{Z}_+$ and $c, d \in \mathbb{C}^r$. Since $\tau = \tau^*$,

taking into account (2.7) and (2.1) and integrating by parts, we find that

$$\begin{aligned}\lambda_j(\Psi(\lambda_j)c|\Psi(\lambda_k)d)_{\mathbb{C}^r} &= \left(\left(\frac{d}{dx} - \tau \right) \Phi(\lambda_j)c|\Psi(\lambda_k)d \right)_{\mathbb{C}^r} \\ &= 2 \left(\varphi(1, \lambda_j, \tau)c|\psi(1, \lambda_k, \tau)d \right)_{\mathbb{C}^r} + \lambda_k \left(\Phi(\lambda_j)c|\Phi(\lambda_k)d \right)_{\mathbb{C}^r}\end{aligned}$$

and, therefore,

$$\lambda_j \Psi^*(\lambda_k) \Psi(\lambda_j) - \lambda_k \Phi^*(\lambda_k) \Phi(\lambda_j) = 2\psi^*(1, \lambda_k, \tau) \varphi(1, \lambda_j, \tau).$$

It follows from the orthogonality of $(\widehat{P}_j)_{j \in \mathbb{Z}_+}$ and $(\widehat{Q}_j)_{j \in \mathbb{N}}$ and relations (2.9) that

$$\alpha_k \Psi_\tau^*(\lambda_k) \Psi_\tau(\lambda_j) \alpha_j = 0 = \alpha_k \Phi_\tau^*(\lambda_k) \Phi_\tau(\lambda_j) \alpha_j, \quad k \neq j.$$

Using this, we obtain the equality

$$\left(\sum_{\lambda_k \in \Delta_n} (-1)^n \alpha_k \psi^*(1, \lambda_k, \tau) \right) \varphi(1, \lambda_j, \tau) \alpha_j = 0, \quad n \in \mathbb{N}, \lambda_j \notin \Delta_n. \quad (4.13)$$

Since (see (2.13))

$$\psi(1, \lambda_k, \tau) = \cos \lambda_k I + \int_0^1 (\cos \lambda_k t) K_{\tau, N}(1, t) dt, \quad k \in \mathbb{N},$$

we can use the Riemann–Lebesgue lemma and the asymptotic behavior of the sequences (λ_k) and (α_k) to prove that

$$\lim_{n \rightarrow \infty} \sum_{\lambda_k \in \Delta_n} (-1)^n \psi(1, \lambda_k, \tau) \alpha_k = \lim_{n \rightarrow \infty} \sum_{\lambda_k \in \Delta_n} \alpha_k = I.$$

Passing to the limit in (4.13) as $n \rightarrow \infty$, we derive the equality $\varphi(1, \lambda_j, \tau) \alpha_j = 0$. Theorem 4.3 is proved. $\square$

4.3. Proof of Proposition 1.3 and Theorems 1.4 and 1.9

Proof of Proposition 1.3. It follows from Theorems 1.1 and 1.2 and Lemma 4.2 proved above that it is sufficient to prove the implication $(\mathbf{a} \in \mathfrak{A}) \implies (\lambda_0 = 0, \alpha_0 > 0)$. This fact, however, is an immediate corollary of Theorem 2.4. $\square$

Proof of Theorem 1.4. (i) Let $\tau_1, \tau \in \text{Re } \mathbf{L}_2$ and $\mathbf{a}_{\tau_1} = \mathbf{a}_\tau = \mathbf{a}$. In view of (2.36) $\mathcal{U}_{\mathbf{a}, \tau_1}^e = \mathcal{I} = \mathcal{U}_{\mathbf{a}, \tau}^e$. We next prove the following implication:

$$\mathcal{U}_{\mathbf{a}, \tau_1}^e = \mathcal{U}_{\mathbf{a}, \tau}^e \implies \mathcal{K}_{\tau_1, N} = \mathcal{K}_{\tau, N}. \quad (4.14)$$

According to (4.1), $\mathcal{U}_{\mathbf{a}, 0}^e = \mathcal{I} + \mathcal{H}_e$, and thus we conclude that the operator $(\mathcal{U}_{\mathbf{a}, 0}^e - \mathcal{I})$ belongs to $\mathcal{B}_2$. Since $\mathcal{U}_{\mathbf{a}, 0}^e > 0$ (see (4.5)), the operator $\mathcal{U}_{\mathbf{a}, 0}^e$ admits a factorization:

$$\mathcal{U}_{\mathbf{a}, 0}^e = (\mathcal{I} + \mathcal{K})^{-1} (\mathcal{I} + \mathcal{K}^*)^{-1}, \quad \mathcal{K} \in \mathcal{B}_2^+.$$

It follows from the equality $\mathcal{U}_{\mathbf{a}, \tau_1}^e = \mathcal{U}_{\mathbf{a}, \tau}^e$ and (2.38) that

$$\begin{aligned}(\mathcal{I} + \mathcal{K}_{\tau_1, N})(\mathcal{I} + \mathcal{K})^{-1} (\mathcal{I} + \mathcal{K}^*)^{-1} (\mathcal{I} + \mathcal{K}_{\tau_1, N}^*) \\ = (\mathcal{I} + \mathcal{K}_{\tau, N})(\mathcal{I} + \mathcal{K})^{-1} (\mathcal{I} + \mathcal{K}^*)^{-1} (\mathcal{I} + \mathcal{K}_{\tau, N}^*).\end{aligned}$$

Therefore, in view of Theorem A.1 we have

$$(\mathcal{I} + \mathcal{K}_{\tau_1, N})(\mathcal{I} + \mathcal{K})^{-1} = (\mathcal{I} + \mathcal{K}_{\tau, N})(\mathcal{I} + \mathcal{K})^{-1},$$

which implies that $\mathcal{K}_{\tau_1, N} = \mathcal{K}_{\tau, N}$.

It follows now from Lemma 2.2 (iii) and (4.14) that $\tau_1 = \tau$.

(ii) Let $q_1, q \in \operatorname{Re} \mathfrak{M}$ and $\mathfrak{b}_{q_1} = \mathfrak{b}_q$. Let us fix $\tau_1 \in b^{-1}(q_1)$ and $\tau \in b^{-1}(q)$. Since $\mathfrak{b}_{q_1} = \mathfrak{b}_q$, the sequences $\mathfrak{a} = \mathfrak{a}_\tau$ and $\mathfrak{a}_1 = \mathfrak{a}_{\tau_1}$ differ at most in the first element. Thus, taking into account (2.36), we obtain that

$$\mathcal{U}_{\mathfrak{a}, \tau}^o = \mathcal{U}_{\mathfrak{a}_\tau, \tau}^o = \mathcal{I} = \mathcal{U}_{\mathfrak{a}_{\tau_1}, \tau_1}^o = \mathcal{U}_{\mathfrak{a}, \tau_1}^o. \quad (4.15)$$

Reasoning as in the proof of (4.14), we establish the implication

$$(\mathcal{U}_{\mathfrak{a}, \tau_1}^o = \mathcal{U}_{\mathfrak{a}, \tau}^o) \implies (\mathcal{K}_{\tau_1, D} = \mathcal{K}_{\tau, D}).$$

Therefore, taking into account (4.15) and the statement (ii) of Lemma 2.2, we get that $b(\tau_1) = b(\tau)$, i.e., $q_1 = q$. The proof is complete. $\square$

Proof of Theorem 1.9. Part (i) is a straightforward consequence of Theorems 4.3 and 1.4. Let us prove part (ii). Assume that $q \in \operatorname{Re} \mathfrak{M}$ and that $\mathfrak{b} = \mathfrak{b}_q$ belongs to $\mathfrak{B}$. Augmenting the sequence $\mathfrak{b}$ to a sequence $\mathfrak{a}$ with $\lambda_0 = 0$ and $\alpha_0 = I$, we get by Proposition 1.3 that $\mathfrak{a} \in \mathfrak{A}$. Set

$$\nu := \nu^{\mathfrak{a}} = I\delta_0 + \mu^{\mathfrak{b}}, \quad \tau := \Theta(H_\nu).$$

From the already proved statement (i) we obtain that $\mathfrak{a} = \mathfrak{a}_\tau$. Then $\mathfrak{b} = \mathfrak{b}_{\tilde{q}}$ for $\tilde{q} = b(\tau)$. It follows from Theorem 1.4 that $q = \tilde{q} = b(\tau)$. The proof is complete. $\square$

Acknowledgments

The authors express their gratitude to DFG for financial support of the project 436 UKR 113/84 and thank the Internationales Zentrum für Komplexe Systeme (IZKS) of Bonn University for the warm hospitality. The research of the first author was partially supported by the Ukrainian DFFD grant F28.1/017.

Appendix A. The spaces

A.1. Sobolev spaces

Suppose that X is a Banach space. Denote by $L_p((a, b), X)$, $1 \leq p < \infty$, the Banach space of all strongly measurable functions $f : (a, b) \rightarrow X$, for which the norm

$$\|f\|_{L_p((a, b), X)} := \left(\int_a^b \|f(t)\|_X^p dt \right)^{1/p}$$

is finite. Denote by $C^k([a, b], X)$ the Banach space of k times continuously differentiable functions $[a, b] \rightarrow X$ with the standard norm.

We also denote by $W_p^1((0, 1), X)$, $1 \leq p < \infty$, the Sobolev space that is the completion of the linear space $C^1([0, 1], X)$ by the norm

$$\|f\|_{W_p^1} := \left(\int_0^1 (\|f(t)\|_X^p dt) \right)^{1/p} + \left(\int_0^1 \|f'(t)\|_X^p dt \right)^{1/p}.$$

Every function $f \in W_p^1((0, 1), X)$ has the derivative f' belonging to $L_p((0, 1), X)$.

We denote by M_r the Banach algebra of $r \times r$ matrices with complex entries. In the standard way the algebra M_r is identified with the Banach algebra of linear operators $A : \mathbb{C}^r \rightarrow \mathbb{C}^r$ and inherits the operator norm

$$\|A\| = \sup_{\|y\|_{\mathbb{C}^r}=1} \|Ay\|_{\mathbb{C}^r},$$

where $\|\cdot\|_{\mathbb{C}^r}$ is the Euclidean norm of $\mathbb{C}^r$ generated by the standard scalar product $(\cdot | \cdot)_{\mathbb{C}^r}$. We shall write L_2 instead of $L_2((0, 1), \mathbb{C}^r)$ and denote by $(\cdot | \cdot)$ the scalar product in L_2 , i.e.,

$$(f | g) = \int_0^1 (f(x) | g(x))_{\mathbb{C}^r} dx, \quad f, g \in L_2.$$

We also use the following notations:

$$\begin{aligned} \mathbf{L}_p &:= L_p((0, 1), M_r), & Re \mathbf{L}_p &:= \{u \in \mathbf{L}_p \mid u = u^*\}, & p \geq 1, \\ W_p^s &:= W_p^s((0, 1), \mathbb{C}^r), & \mathbf{W}_p^s &:= W_p^s((0, 1), M_r) & (s \in \mathbb{Z}, p \geq 1). \end{aligned}$$

Set $W_{2,0}^1(0, 1) = \{f \in W_2^1(0, 1) \mid f(0) = f(1) = 0\}$. Recall (see [39]) that $W_2^{-1}(0, 1)$ is the dual space of $W_{2,0}^1(0, 1)$. Denote by $\mathbf{W}_2^{-1}$ the Banach space of matrices $f = (f_{ij})_{1 \leq i, j \leq r}$ with entries $f_{ij} \in W_2^{-1}(0, 1)$ and with the inherited norm

$$\|f\|_{\mathbf{W}_2^{-1}} = \left(\sum_{1 \leq i, j \leq r} \|f_{ij}\|_{W_2^{-1}(0, 1)}^2 \right)^{1/2}.$$

Let $Re \mathbf{W}_2^{-1} := \{f \in \mathbf{W}_2^{-1} \mid f = f^*\}$. Here $f^* := (\overline{f_{ji}})_{1 \leq i, j \leq r}$ for $f = (f_{ij})_{1 \leq i, j \leq r}$.

A.2. Factorization of operators

We state below some necessary facts related to the factorization of Fredholm operators [36, Ch.4]. Denote by $\mathcal{B}_2$ the Hilbert space of all Hilbert–Schmidt operators in L_2 . Every operator $\mathcal{K} \in \mathcal{B}_2$ is an integral operator with kernel $K \in L_2((0, 1)^2, M_r)$. Set $\Omega := \{(x, t) \mid 0 \leq t \leq x \leq 1\}$, $\Omega^- := [0, 1]^2 \setminus \Omega$ and denote by $\mathcal{B}_2^+$ and $\mathcal{B}_2^-$ the subalgebras of $\mathcal{B}_2$ consisting of all operators with kernels K satisfying the condition $K(x, t) = 0$ a.e. in Ω^- and $K(x, t) = 0$ a.e. in Ω , respectively. It is obvious that $\mathcal{B}_2 = \mathcal{B}_2^+ \oplus \mathcal{B}_2^-$; moreover, the operators in $\mathcal{B}_2^\pm$ are Volterra ones [36, Ch. IV]. Denote by $\mathcal{P}^+$ an orthogonal projector in $\mathcal{B}_2$ onto $\mathcal{B}_2^+$.

We say that an operator $\mathcal{I} + \mathcal{K}$ with $\mathcal{K} \in \mathcal{B}_2$ admits a factorization in $\mathcal{B}_2$ if

$$\mathcal{I} + \mathcal{K} = (\mathcal{I} + \mathcal{K}_+)^{-1}(\mathcal{I} + \mathcal{K}_-)^{-1} \tag{A.1}$$

with some $\mathcal{K}_+ \in \mathcal{B}_2^+$ and $\mathcal{K}_- \in \mathcal{B}_2^-$.

Theorem A.1 Assume that $\mathcal{K} \in \mathcal{B}_2$ with kernel K is such that $\mathcal{I} + \mathcal{K}$ admits a factorization in $\mathcal{B}_2$. Then the operators $\mathcal{K}_+$ and $\mathcal{K}_-$ in (A.1) are determined uniquely by $\mathcal{K}$. Moreover, the operators $\mathcal{K}_+$ and $\mathcal{K}_-^*$ are solutions for $\mathcal{X} \in \mathcal{B}_2$ of the equations

$$\mathcal{X} + \mathcal{P}^+ \mathcal{K} + \mathcal{P}^+ \mathcal{X} \mathcal{K} = 0, \quad \mathcal{X} + \mathcal{P}^+ \mathcal{K}^* + \mathcal{P}^+ \mathcal{X} \mathcal{K}^* = 0 \quad (\text{A.2})$$

respectively. For the operator $\mathcal{I} + \mathcal{K}$ to admit a factorization in $\mathcal{B}_2$, it is necessary and sufficient that one of the following conditions should hold:

(i) for every $a \in [0, 1]$ the integral equation

$$f(x) + \int_0^a K(x, t) f(t) dt = 0, \quad x \in (0, 1),$$

has only the trivial solution in L_2 ;

(ii) at least one of the equations in (A.2) has a solution.

In terms of the kernels the first equation in (A.2) takes the form

$$X(s, t) + K(s, t) + \int_0^s X(s, \xi) K(\xi, t) d\xi = 0, \quad (s, t) \in \Omega. \quad (\text{A.3})$$

Theorem A.1 yields the following connection between the Krein accelerants and the Krein equation, namely:

Proposition A.2 Assume that $H \in L_2((-1, 1), M_r)$ and that $\mathcal{H}$ is an integral operator of (3.1). Then the following statements are equivalent:

- (i) the function H belongs to the class $\mathfrak{H}_2$;
- (ii) the operator $\mathcal{I} + \mathcal{H}$ admits a factorization;
- (iii) the Krein equation (1.11) has a solution in $L_2(\Omega, M_r)$.

Theorem A.1 also implies the following lemma.

Lemma A.3 If K belongs to $L_2((0, 1)^2, M_r)$, then equation (A.3) has at most one solution, and the equation

$$X(s, t) + \int_0^s X(s, \xi) K(\xi, t) dt = 0, \quad (s, t) \in \Omega,$$

has only zero solution in $L_2(\Omega, M_r)$.

We denote by G_2 the set of all functions $K : [0, 1]^2 \rightarrow M_r$ having the property that the mappings

$$[0, 1] \ni x \mapsto K(x, \cdot) \in \mathbf{L}_2, \quad [0, 1] \ni t \mapsto K(\cdot, t) \in \mathbf{L}_2$$

are continuous on the interval $[0, 1]$. It is obvious that $G_2 \subset L_2((0, 1)^2, M_r)$. The set G_2 becomes a Banach space upon introducing the norm

$$\|K\|_{G_2} := \max \left\{ \max_{x \in [0, 1]} \|K(x, \cdot)\|_{\mathbf{L}_2}, \max_{t \in [0, 1]} \|K(\cdot, t)\|_{\mathbf{L}_2} \right\}.$$

Denote by G_2^+ and G_2^- the subspaces of G_2 consisting of all functions K that satisfy the condition $K(x, t) = 0$ a.e. in Ω^- and $K(x, t) = 0$ a.e. in Ω , respectively.

Also denote by $\mathfrak{G}_2$ a Banach algebra consisting of integral operators $\mathcal{K}$ with kernels K belonging to G_2 . The norm in $\mathfrak{G}_2$ is defined via

$$\|\mathcal{K}\|_{\mathfrak{G}_2} = \|K\|_{G_2}, \quad \mathcal{K} \in \mathfrak{G}_2.$$

Denote by $\mathfrak{G}_2^+$ and $\mathfrak{G}_2^-$ the set of those $\mathcal{K} \in \mathfrak{G}_2$, whose kernels belong respectively to G_2^+ and G_2^- . Observe that $\mathfrak{G}_2^+$ and $\mathfrak{G}_2^-$ form closed subalgebras of $\mathfrak{G}_2$ and that $\mathfrak{G}_2^\pm \subset \mathcal{B}_2^\pm$.

Lemma A.4 *The mapping $A \mapsto (\mathcal{I} + A)^{-1} - \mathcal{I} =: \eta(A)$ acts continuously in G_2^+ .*

Proof. Using the Cauchy–Bunyakovski inequality, we find that for every $A \in G_2^+$

$$\|A^{n+1}\|_{G_2^+} \leq (n!)^{-1/2} \|A\|_{G_2^+}^{n+1}, \quad n \in \mathbb{Z}_+. \quad (\text{A.4})$$

Let $A, B \in G_2^+$ and $\|A\|_{G_2^+}, \|B\|_{G_2^+} \leq r$ for some fixed $r > 0$. Since

$$\eta(A) - \eta(B) = \sum_{n=1}^{\infty} [(-A)^n - (-B)^n],$$

in view of (A.4) and the identity $A^{n+1} - B^{n+1} = \sum_{k=0}^n A^{n-k}(A - B)B^k$, straightforward calculations show that the mapping η acts continuously in G_2^+ and that

$$\|\eta(A) - \eta(B)\|_{G_2^+} \leq C^2(r) \|A - B\|_{G_2^+}$$

with

$$C(r) := 1 + \sum_{k=0}^{\infty} (k!)^{-1/2} r^{k+1}.$$

The proof is complete. $\square$

Appendix B. Lemmas about orthogonal projectors

Suppose that H is a Hilbert space and denote by $\mathcal{B}(H)$ the Banach algebra of all bounded linear operators in H .

Lemma B.1 *Let $(P_n)_{n=1}^\infty$ and $(G_n)_{n=1}^\infty$ be sequences of pairwise orthogonal projectors in H of finite rank such that $\sum_{n=1}^\infty P_n = \sum_{n=1}^\infty G_n = I_H$ and $\sum_{n=1}^\infty \|P_n - G_n\|^2 < \infty$. Then there exists $N_0 \in \mathbb{N}$ such that, for all $N \geq N_0$,*

$$\sum_{n=1}^N \text{rank } P_n = \sum_{n=1}^N \text{rank } G_n.$$

Proof. Let us choose a natural N_0 such that $\sum_{n=N_0}^\infty \|P_n - G_n\|^2 < \frac{1}{4}$. Fix an arbitrary natural $N \geq N_0$ and consider the orthogonal projectors $P = \sum_{n=1}^N P_n$ and $G = \sum_{n=1}^N G_n$. We shall now show that $\|P - G\| < 1$. Indeed, the assumptions of the lemma imply that

$$P - G = \sum_{n=N+1}^{\infty} (P_n - G_n) = \sum_{n=N+1}^{\infty} P_n(P_n - G_n) + \sum_{n=N+1}^{\infty} (P_n - G_n)G_n.$$

Thus for every $f \in H$ of norm 1 we get the inequality

$$|((P - G)f|f)_H| \leq \sum_{n=N+1}^{\infty} (\|P_n f\| + \|G_n f\|) \|(P_n - G_n)f\|.$$

Using the Cauchy–Bunyakovski inequality, we obtain that

$$\|P - G\|^2 \leq 4 \sum_{n=N+1}^{\infty} \|P_n - G_n\|^2 < 1.$$

The inequality $\|P - G\| < 1$ implies (see [40, Ch.1]) that $\text{rank } P = \text{rank } G$, and the proof is complete. $\square$

Lemma B.2 *Assume that $(A_j)_{j=1}^{\infty}$ is a sequence in $\mathcal{B}(H)$ and that $(G_j)_{j=1}^{\infty}$ is a sequence of pairwise orthogonal projectors such that the following holds:*

- (i) *the series $\sum_{j=1}^{\infty} A_j$ converges in the strong operator topology to an operator A ;*
- (ii) *the orthogonal projector $G := I_H - \sum_{j=1}^{\infty} G_j$ is of finite rank;*
- (iii) *$\sum_{j=1}^{\infty} \|A_j - G_j\|^2 < 1$ and $\text{rank } A_j \leq \text{rank } G_j < \infty$ for every $j \in \mathbb{N}$.*

Then $\text{codim } \text{Ran } A \geq \text{rank } G$.

Proof. It follows from the assumptions of the lemma that, for every $f \in L_2$,

$$\sum_{j=1}^{\infty} \|(A_j G_j - G_j)f\| \leq \sum_{j=1}^{\infty} \|A_j - G_j\| \|G_j f\| \leq \left(\sum_{j=1}^{\infty} \|A_j - G_j\|^2 \right)^{1/2} \|f\|$$

and, therefore, the series $\sum_{j=1}^{\infty} (A_j G_j - G_j)$ strongly converges to some operator $B \in \mathcal{B}(H)$ with $\|B\| < 1$. Hence the series $\sum_{j=1}^{\infty} A_j G_j$ strongly converges to the operator $\tilde{A} := (I_H + B)(I_H - G)$. Since the operator $I_H + B$ is invertible, $\tilde{A}$ has a closed range and $\text{codim } \text{Ran } \tilde{A} = \text{rank } G$. Since $\text{rank } A_j \leq \text{rank } G_j < \infty$, we have

$$\text{Ran } A_j G_j = \text{Ran } A_j \tag{B.1}$$

for all $j \in \mathbb{N}$. Indeed, if (B.1) does not hold for some $k \in \mathbb{N}$, then there exists $c \in \text{Ran } G_k$ such that $A_k c = 0$. Therefore $\|(A_k - G_k)c\| = \|c\|$, which implies that $\|A_k - G_k\| \geq 1$ thus contradicting assumption (iii).

We denote by X the closed linear hull of the set $\cup_{j \in \mathbb{N}} \text{Ran } A_j$. It is obvious that $\text{Ran } A \subset X$. It follows from (B.1) and the definition of the operator $\tilde{A}$ that $\text{Ran } \tilde{A}$ contains the set $\cup_{j \in \mathbb{N}} \text{Ran } A_j$. But, as already noted, $\text{Ran } \tilde{A}$ is a closed set. Thus $\text{Ran } A \subset X \subset \text{Ran } \tilde{A}$ and, therefore, $\text{codim } \text{Ran } A \geq \text{codim } \text{Ran } \tilde{A} = \text{rank } G$. The proof is complete. $\square$

Lemma B.3 *Let $\{A_j\}_{j=0}^n$ be a set of self-adjoint operators from the algebra $\mathcal{B}(H)$ that are of finite rank for $j \neq 0$. If*

$$\sum_{j=0}^n A_j = I_H, \quad \sum_{j=1}^n \text{rank } A_j \leq \text{codim } \text{Ran } A_0,$$

then $\{A_j\}_{j=0}^n$ is the set of pairwise orthogonal projectors.

Proof. It follows from the assumptions of the lemma that the space H is the direct sum of the subspaces $\text{Ran } A_j$ for $j = 0, \dots, n$ and that $A_k(A_k - I_H) + \sum_{j \neq k} A_j A_k = 0$ for every $k \leq n$. It is obvious that the equality is possible only if all the summands on the left hand side are equal to zero. This immediately yields the statement of the lemma. $\square$

References

- [1] Marchenko V A 1977 *Sturm–Liouville Operators and their Applications* (Kiev: Naukova Dumka) (in Russian) (Engl. transl. 1986 (Basel: Birkhäuser))
- [2] Levitan B M 1987 *Inverse Sturm–Liouville Problems* (Utrecht: VNU Science Press)
- [3] Pöschel J and Trubowitz E 1987 *Inverse Spectral Theory (Pure Appl. Math. vol 130)* (Boston, MA: Academic Press, Inc.)
- [4] Gesztesy F 2007 Inverse spectral theory as influenced by Barry Simon in: *Spectral Theory and Mathematical Physics: A Festschrift in Honor of Barry Simon’s 60th Birthday, Part 2*, in: *Proc. Sympos. Pure Math.* **76** (Providence, RI: American Mathematical Society) pp 741–820
- [5] Borg G 1946 Eine Umkehrung der Sturm–Liouvilleschen Eigenwertaufgabe. Bestimmung der Differentialgleichung durch die Eigenwerte (*Acta Math.* 78) pp 1–96 (in German)
- [6] Marchenko V A 1950 Concerning the theory of a differential operator of the second order *Dokl. Akad. Nauk SSSR* **72** 457–460 (in Russian)
- [7] Gelfand I M and Levitan B M 1951 On the determination of a differential equation by its spectral function *Izv. Akad. Nauk SSSR Ser. Mat.* **15** 309–60 (in Russian)
- [8] Marchenko V A 1952 Some questions in the theory of one-dimensional linear differential operators of the second order **I**, *Trudy Moskov. Mat. Ob-va* **1** 327–420 (in Russian) (Engl. transl. 1973 *Amer. Math. Soc.* **101** 1–104)
- [9] Krein M G 1951 Solution of the inverse Sturm–Liouville problem *Dokl. Akad. Nauk SSSR* **76** 21–4 (in Russian)
- [10] Krein M G 1953 On the transfer function of a one-dimensional boundary problem of the second order *Dokl. Akad. Nauk SSSR* **88** 405–8 (in Russian)
- [11] Krein M G 1954 On a method of effective solution of an inverse boundary problem *Dokl. Akad. Nauk SSSR* **94** 987–90 (in Russian)
- [12] Krein M G 1954 On integral equations generating differential equations of 2nd order *Dokl. Akad. Nauk SSSR* **97** no 1 21–4
- [13] Ramm A G 2000 Krein’s method in inverse scattering In: *Operator theory and Applications*, in: *Fields Institute Commun.* vol 25 (Providence: American Mathematical Society) pp 441–56
- [14] Isaacson E and Trubowitz E 1983 The inverse Sturm–Liouville problem: I *Commun. Pure Appl. Math.* **36** no 6 767–83
- [15] Isaacson E, McKean H and Trubowitz E 1984 The inverse Sturm–Liouville problem: II *Commun. Pure Appl. Math.* **37** no 1 1–11
- [16] Dahlberg B E J and Trubowitz J E 1984 The inverse Sturm–Liouville problem: III *Commun. Pure Appl. Math.* **37** no 2 255–67
- [17] Chelkak D, Kargaev P and Korotyaev E 2004 Inverse problem for harmonic oscillator perturbed by potential, characterization *Commun. Math. Phys.* **249** no 1 133–96
- [18] Chelkak D and Korotyaev E 2007 The inverse problem for perturbed harmonic oscillator on the half-line with a Dirichlet boundary condition *Ann. Henri Poincaré* **8** no 6 1115–50
- [19] Newton R G and Jost R 1955 The construction of potentials from S–matrix for systems of differential equations *Nuovo Cimento* **1** 590–622
- [20] Krein M G 1955 On determination of the potential of a particle from its S-function *Dokl. Akad. Nauk SSSR* **105** no 3 433–6

- [21] Agranovich Z S and Marchenko V A 1963 *The Inverse Problem of Scattering Theory Translated from the Russian by B. D. Seckler Gordon and Breach Science Publishers* (New York-London) pp xiii+291
- [22] Gasymov M G and Levitan B M 1966 Inverse problem for the Dirac system *Dokl. Akad. Nauk SSSR* **167** 967–70 (in Russian)
- [23] Gasymov M G 1968 An inverse problem of scattering theory for a system of Dirac equations of order $2n$ *Trudy Moskov. Mat. Ob-va* **19** 41–112 (in Russian)
- [24] Lesch M and Malamud M M The inverse spectral problem for first order systems on the half line in: *Differential operators and related topics* Vol. I (Odessa, 1997), 199–238, in: *Oper. Theory Adv. Appl.* vol **117** Birkhäuser, Basel, 2000.
- [25] Carlson R 2002 An inverse problem for the matrix Schrödinger equation *Math. Anal. Appl.* **267** no 2 564–75
- [26] Yurko V A 2006 Inverse problems for matrix Sturm–Liouville operators *Russian J. Math. Phys.* **13** no 1 111–8
- [27] Malamud M M 2005 Uniqueness of the matrix Sturm–Liouville equation given a part of the monodromy matrix, and Borg type results *Sturm–Liouville theory* (Basel: Birkhäuser) pp 237–70
- [28] Yurko V A 2006 Inverse problems for the matrix Sturm–Liouville equation on a finite interval *Inverse Problems* **22** 1139–49
- [29] Chelkak D and Korotyaev E 2006 Parametrization of the isospectral set for the vector-valued Sturm–Liouville problem *J. Funct. Anal.* **241** no 1 359–73
- [30] Chelkak D and Korotyaev E 2009 Weyl–Titchmarsh functions of vector-valued Sturm–Liouville operators on the unit interval *J. Funct. Anal.* doi:10.1016/j.jfa.2009.05.010
- [31] Kappeler T, Perry P, Shubin M and Topalov P 2005 The Miura map on the line *Intern. Math. Research Notes* **50** 3091–133
- [32] Krein M G 1956 Towards the theory of accelerants and S–matrices of canonical differential systems *Dokl. Akad. Nauk SSSR* **111** 1167–70 (in Russian)
- [33] Albeverio S, Hryniv R and Mykytyuk Ya 2005 Inverse spectral problems for Dirac operators with summable potentials *Russian J. Math. Phys.* **12** 406–23
- [34] Albeverio S, Hryniv R and Mykytyuk Ya 2005 Inverse spectral problems for Sturm–Liouville operators in impedance form *J. Funct. Anal.* **222** 143–77
- [35] Trush N S 2008 Solutions of the Cauchy problem for factorized Sturm–Liouville equation in a Banach algebra *Mat. Stud.* **31** 75–82
- [36] Gohberg I and Krein M 1967 *Theory of Volterra Operators in Hilbert Space and its Applications* (Moscow: Nauka) (in Russian) (Engl. transl. 1970 *Amer. Math. Soc. Transl. Math. Monographs* **24** (Providence, RI: American Mathematical Society))
- [37] Titchmarsh E C 1937 *Introduction to the Theory of Fourier Integrals* (Oxford University Press)
- [38] Trush N S 2008 Asymptotics of singular values of entire matrix-valued sine-type functions *Mat. Stud.* **30** 95–7
- [39] Adams R A and Fournier J F 2003 *Sobolev Spaces*. 2nd ed (*Pure and Applied Mathematics* **140**)(New York: Academic Press)
- [40] Kato T 1995 *Perturbation Theory for Linear Operators* (Springer–Verlag: New York)